

HEMOLAX FILM-COATED TABLET

Diosmin/Hesperidin (450 mg/50 mg)

What is in this leaflet

1. What Hemolax Film-Coated Tablet is used for
2. How Hemolax Film-Coated Tablet works
3. Before you use Hemolax Film-Coated Tablet
4. How to use Hemolax Film-Coated Tablet
5. While you are using Hemolax Film-Coated Tablet
6. Side effects
7. Storage and Disposal of Hemolax Film-Coated Tablet
8. Product Description
9. Manufacturer
10. Product Registration Holder
11. Date of revision
12. Serial Number

What Hemolax Film-Coated Tablet is used for

It is recommended for the treatment for venous circulation disorders (heavy legs, pain, early morning restless legs) and the treatment of symptoms related to hemorrhoids (acute hemorrhoidal attack).

How Hemolax Film-Coated Tablet works

This medicine strengthens veins by increasing venous tone, and protects small blood vessels.

Before you use Hemolax Film-Coated Tablet

When you must not use it

Do not take Hemolax Film-Coated Tablet:

If you are allergic (hypersensitive) to diosmin, hesperidin, micronized purified flavonoid fraction or any of the other ingredients of this medicine (see Ingredients section).

Do not use in children below 18 years old unless advised by your doctor.

Before you start to use it

Ask your doctor or pharmacist for advice before taking this medicine.

Pregnancy and breast-feeding:

If you are pregnant, think you may be pregnant, planning to have a baby, or breast-feeding, ask your doctor or pharmacist for advice before taking this medicine.

Taking other medicines

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

How to use Hemolax Film-Coated Tablet

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

How much to use & When to use it

Venous circulation disorders

The recommended dose is 2 tablets daily; i.e. one at midday and one in the evening at mealtimes.

Acute hemorrhoidal attack

The recommended dose is 6 tablets daily (in 3 divided doses) for the first 4 days, then 4 tablets daily (in 2 divided doses) for the next 3 days.

How long to use it

Continue taking Hemolax Film-Coated Tablet for as long as your doctor or pharmacist recommends.

If you forget to use it

Do not take a double dose to make up for a forgotten dose.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

If you use too much (overdose)

If you take more Hemolax Film-Coated Tablet than you should, contact your doctor or pharmacist immediately.

Symptoms which may be experienced with overdose of Hemolax Film-Coated Tablet include diarrhoea, nausea, abdominal pain, itching and rash.

While you are using it

Things you must do

Take your medicine exactly as your doctor or pharmacist has told you.

Tell all the doctors, dentists and pharmacists treating you that you are taking Hemolax Film-Coated Tablet.

Things you must not do

This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.

Side effects

Like all medicines, Hemolax Film-Coated Tablet can cause side effects, although not everybody gets them.

They may include:

Diarrhoea, dyspepsia (difficult digestion), nausea, vomiting, colitis (inflammation of the large intestine), dizziness, headache, discomfort, rash (sudden redness in patches), pruritus (skin reactions such as itching) and urticaria (allergic reaction in the form of small red spots), abdominal pain, isolated oedema (swelling of the face, lips and eyelids), Quincke's oedema (swelling of the face, lips, mouth, tongue or throat which could cause breathing difficulties).

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. By reporting side effects, you can help provide more information on the safety of this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npa.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

HEMOLAX FILM-COATED TABLET

Diosmin/Hesperidin (450 mg/50 mg)

Storage and Disposal of Hemolax Film-Coated Tablet**Storage**

The expiry date of this pack is printed on the box. Do not use this pack after this date. Keep all medicines out of reach of children.

Do not store above 30°C.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help protect the environment.

Product description**What it looks like**

Hemolax Film-Coated Tablet is brown-yellow, oval-shaped film-coated tablet, biconvex and plain on both sides.

Ingredients**Active ingredient:**

Each film-coated tablet contains:

Micronized purified flavonoid fraction corresponding to:

Diosmin..... 450 mg

Flavonoids expressed as hesperidin ..50 mg

The other ingredients are:

Microcrystalline cellulose, sodium starch glycolate, magnesium stearate, gelatin, hypromellose, macrogol 6000, talc, titanium dioxide, sodium lauryl sulfate, yellow iron oxide, red iron oxide, glycerin.

MAL No.:

Manufacturer:

STELLAPHARM J.V. CO., LTD. - BRANCH 1

(A Joint Venture Company of Auxilto GmbH, Germany)

No. 40 Tu Do Avenue, Vietnam-Singapore Industrial Park, An Phu Ward, Thuan An Town, Binh Duong Province, Vietnam.

Product Registration Holder

Stadpharm Sdn. Bhd.

E-9-01, Oasis Square, Ara Damansara, No. 2, Jalan PJU 1A/7A, 47301 Petaling Jaya, Selangor Darul Ehsan, Malaysia.

Date of revision:

18 July, 2022

Serial Number

NPRA (R2) 21/193