

PARTAMOL TAB. TABLETS 500MG

Paracetamol 500 mg

What is in this leaflet

1. What PARTAMOL TAB. TABLETS 500MG is used for
2. How PARTAMOL TAB. TABLETS 500MG works
3. Before you use PARTAMOL TAB. TABLETS 500MG
4. How to use PARTAMOL TAB. TABLETS 500MG
5. While you are using PARTAMOL TAB. TABLETS 500MG
6. Side effects
7. Storage and Disposal of PARTAMOL TAB. TABLETS 500MG
8. Product Description
9. Manufacturer
10. Product Registration Holder
11. Date of Revision
12. Serial Number

1. What PARTAMOL TAB. TABLETS 500MG is used for

Paracetamol is a mild analgesic and antipyretic, and is recommended for the treatment of most painful and febrile conditions, for example, headache including migraine and tension headaches, toothache, backache, rheumatic and muscle pains, dysmenorrhoea, sore throat, and for relieving the fever, aches and pains of colds and flu. Also recommended for the symptomatic relief of pain due to non-serious arthritis.

2. How PARTAMOL TAB. TABLETS 500MG works

Paracetamol is an antipyretic analgesic. The mechanism of action is probably similar to that of aspirin and dependant on the inhibition of prostaglandin synthesis. This inhibition appears, however to be on a selective basis.

3. Before you use PARTAMOL TAB. TABLETS 500MG

- When you must not use it

Do not use if you are allergic to paracetamol or any of the other ingredients in the product.

Do not use this medicine if you are taking any other prescription or non-

prescription medicines containing paracetamol to treat pain, fever, symptoms of cold and flu, or to aid sleep.

Do not take PARTAMOL TAB. TABLETS 500MG if you are pregnant, trying to get pregnant or think you may be pregnant.

Ask your doctor or pharmacist for advice before taking any medicine.

- Before you start to use it

Check with your doctor before use if:

- you have liver or kidney problems.
- are underweight or malnourished.
- regularly drink alcohol.

You may need to avoid using this product altogether or limit the amount of paracetamol that you take.

Check with your doctor before use if:

- you have a severe infection as this may increase the risk of metabolic acidosis. Signs of metabolic acidosis include: deep, rapid, difficult breathing, feeling sick (nausea), being sick (vomiting), loss of appetite. Contact a doctor immediately if you get a combination of these symptoms.

Please see your doctor if your symptoms do not improve.

You can take this product with stomach ulcer.

You can take this product whilst breast feeding.

- Taking other medicines

Check with your doctor before use if you are taking warfarin or similar medicines used to thin the blood.

Always read and follow the label.

Keep medicine out of sight and reach of children.

Maximum duration of continued use without medical advice: 3days.

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop. Some medications and this product can interfere with each other.

4. How to use PARTAMOL TAB. TABLETS 500MG

- How much to use

- Adults (including the elderly) and children aged 12 years and over: 1 to 2 tablets taken every 4 to 6 hours as required.

Maximum daily dose: 4000mg.

- Children, 6 to 11 years: ½ - 1 tablet every 4 to 6 hours as required.

Maximum daily dose: 2000mg.

Do not exceed the stated dose.

The lowest dose necessary to achieve efficacy should be used.

Do not take more frequently than every 4 hours.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Taken every 4 to 6 hours as required.

Adults (including the elderly) and children aged 12 years and over: Maximum daily dose 4000mg (8 tablets in 24 hours).

Children 6 to 11 years: Maximum daily dose 2000mg (4 tablets in 24 hours).

Minimum dosing interval: 4hours.

Use as directed by your doctor or pharmacist.

- How long to use it

Continue taking PARTAMOL TAB. TABLETS 500MG for as long as your doctor recommends.

If symptoms persist for more than 3 days, please see your doctor

- If you forget to use it

Consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

If you take more of the medicine than you should seek medical advice immediately because of the risk of liver failure. Paracetamol overdose may cause liver failure. Contact your doctor immediately or go to the Emergency

PARTAMOL TAB. TABLETS 500MG

Paracetamol 500 mg

Department of your nearest hospital, if you think you or anyone else may have taken too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

5. While you are using PARTAMOL TAB. TABLETS 500MG

- Things you must do

Take your medicine exactly as your doctor has told you or according to instruction on pack.

Tell all the doctors, dentists and pharmacists treating you that you are taking PARTAMOL TAB. TABLETS 500MG.

Tell your doctor immediately if you become pregnant while taking this medication.

- Things you must not do

Do not exceed the stated dose.

Do not take more than 4000mg paracetamol per day (adults and children over 12 years).

Do not take more than 2000mg paracetamol per day (children, 6 to 11 years).

Not recommended for children under the age of 6 years.

Do not take for more than 3 days without asking your doctor.

Do not stop taking the medicine unless advised by your doctor.

Do not take any new medicines without consulting your doctor.

Do not give PARTAMOL TAB. TABLETS 500MG to anyone else, even if they have the same symptoms or condition as you.

Do not take with any other product containing paracetamol.

- Things to be careful of

This preparation contains PARACETAMOL. Do not take any other paracetamol containing medicines at the same time.

ALLERGY ALERT: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters and rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

6. Side effects

Stop taking this medicine and tell your doctor immediately if:

- you experience allergic reactions such as skin rash or itching, sometimes with breathing problems or swelling of the lips, tongue, throat or face.
- you experience a skin rash or peeling, or mouth ulcers.
- you have previously experienced breathing problems with aspirin or non-steroidal anti-inflammatories, and experience a similar reaction with this product.
- you experience unexplained bruising or bleeding

These reactions are rare.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

7. Storage and Disposal of PARTAMOL TAB. TABLETS 500MG

- Storage

Keep out of reach of children.

Store in a well-closed container, in a dry place, protect from light. Do not store above 30°C.

Do not use PARTAMOL TAB. TABLETS 500MG after the expiry date which is stated on the box.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

8. Product description

- What it looks like

White, round-shaped tablet, engraved "PARACETAMOL" on one side and breakable line on the other side.

- Ingredients

- Active ingredient:

Paracetamol

- Inactive ingredient(s):

Microcrystalline cellulose, Sodium starch glycolate, Povidone K90, Magnesium stearate.

- MAL number:

MALXXXXX

9. Manufacturer:

STELLAPHARM J.V. CO., LTD. - BRANCH 1

(A Joint Venture Company of Auxilto GmbH, Germany)

No. 40 Tu Do Avenue, Vietnam-Singapore Industrial Park, An Phu Ward, Thuan An Town, Binh Duong Province, Vietnam.

10. Product Registration Holder

Stadpharm Sdn. Bhd.

E-9-01, Oasis Square, Ara Damansara, No. 2, Jalan PJU 1A/7A, 47301 Petaling Jaya, Selangor Darul Ehsan, Malaysia.

11. Date of revision:

15/07/2024

12. Serial Number

NPRA (R2) 24/097