HONG KONG

multiBic® 2 mmol/L potassium
Solution for haemofiltration
For intravenous use only

Double chamber bag

1000ml of the ready-to-use solution contains:

Sodium chloride	6.136 g
Sodium hydrogen carbonate	2.940 g
Potassium chloride	0.1491 g
Calcium chloride dihydrate	0.2205 g
Magnesium chloride hexahydrate	0.1017 g
Glucose monohydrate	1.100 g
(anhydrous glucose	1.000 g)

Water for injections, hydrochloric acid 25%, carbon dioxide

Sterile and free from bacterial endotoxins. Do not use unless the solution is clear and colourless and the container is undamaged. For single use only; any unused solution must be discarded. **Store below 30°C. Do not store below 4°C.** Remove the overwrap immediately before use. Do not use before the two solutions have been thoroughly mixed. The ready-to-use solution should be used immediately. However if this is not possible the solution must be used within 48 hours after mixing. Instructions for use – please refer to the Patient Information Leaflet. Please read the leaflet carefully before using this solution. Keep out of the reach and sight of children. To be used only as directed by a medical practitioner.
請遵照醫生指示使用，請將產品儲存於溫度攝氏30°C以下及4°C以上。

HK-XXXXX

INDIA

multiBic® 2 mmol/L potassium
Solution for haemofiltration
For intravenous use only

Double chamber bag

1000ml of the ready-to-use solution contains:

Sodium chloride	6.136 g
Sodium hydrogen carbonate	2.940 g
Potassium chloride	0.1491 g
Calcium chloride dihydrate	0.2205 g
Magnesium chloride hexahydrate	0.1017 g
Glucose monohydrate	1.100 g
(anhydrous glucose	1.000 g)

Water for injections, hydrochloric acid 25%, carbon dioxide

Sterile and free from bacterial endotoxins. Do not use unless the solution is clear and colourless and the container is undamaged. To be used as directed by your doctor. For single use only; any unused solution must be discarded. **Store below 30°C. Do not store below 4°C.** Remove the overwrap immediately before use. Do not use before the two solutions have been thoroughly mixed. The ready-to-use solution should be used immediately. However if this is not possible the solution must be used within 48 hours after mixing. Instructions for use – please refer to the Patient Information Leaflet. Please read the leaflet carefully before using this solution. Keep out of the reach and sight of children.
Imported by: Fresenius Medical Care (I) Pvt. Ltd.,
B-7 & B-8, Middle Portion, SIPCOT Industrial Park, Irungattukottai, Sriperumbudur-602105, Chennai, INDIA
Marketed by: Fresenius Medical Care (I) Pvt. Ltd.
S-21, Star City, District Centre Mayur Place, Mayur Vihar Phase - 1, New Delhi – 110091, INDIA
Import Licence no.: MRP (inclusive of tax):
Multi-lingual, multi-country label

PHILIPPINES
Haemofiltration Solution

multiBic® 2 mmol/L potassium
Solution for haemofiltration
For intravenous use only

Double chamber bag

1000ml of the ready-to-use solution contains:

Sodium chloride	6.136 g
Sodium hydrogen carbonate	2.940 g
Potassium chloride	0.1491 g
Calcium chloride dihydrate	0.2205 g
Magnesium chloride hexahydrate	0.1017 g
Dextrose	1.100 g
(anhydrous glucose	1.000 g)

Water for injections, hydrochloric acid 25%, carbon dioxide

Sterile and free from bacterial endotoxins (non-pyrogenic). Do not use unless the solution is clear and colourless and the container is undamaged. To be used as directed by your doctor. For single use only; any unused solution must be discarded. **Store below 30°C. Do not store below 4°C.** Remove the overwrap immediately before use. Do not use before the two solutions have been thoroughly mixed. The ready-to-use solution should be used immediately. However if this is not possible the solution must be used within 48 hours after mixing. Instructions for use – please refer to the Patient Information Leaflet. Please read the leaflet carefully before using this solution. Keep out of the reach and sight of children.
Imported and Distributed by: Fresenius Medical Care Philippines, Inc., 2257 Don Chino Roces Ave., Makati City, 1231 Philippines
Caution: Food, Drugs, Devices, and Cosmetics Act prohibits dispensing without prescription.
DR – XY0000

AUSTRALIA / MALAYSIA

multiBic® 2 mmol/L potassium
Solution for haemofiltration
For intravenous use only

Double chamber bag

1000ml of the ready-to-use solution contains:

Sodium chloride	6.136 g
Sodium hydrogen carbonate	2.940 g
Potassium chloride	0.1491 g
Calcium chloride dihydrate	0.2205 g
Magnesium chloride hexahydrate	0.1017 g
Glucose monohydrate	1.100 g
(anhydrous glucose	1.000 g)

Water for injections, hydrochloric acid 25%, carbon dioxide

Sterile and free from bacterial endotoxins. Do not use unless the solution is clear and colourless and the container is undamaged. To be used as directed by your doctor. For single use only; any unused solution must be discarded. **Store below 30°C. Do not store below 4°C.** Remove the overwrap immediately before use. Do not use before the two solutions have been thoroughly mixed. The ready-to-use solution should be used immediately. However if this is not possible the solution must be used within 48 hours after mixing. Instructions for use – please refer to the Patient Information Leaflet/ Package Insert. Please read the leaflet/insert carefully before using this solution. Keep out of the reach and sight of children.
Distributed by: Fresenius Medical Care Australia Pty. Ltd.
305 Woodpark Road, Smithfield NSW 2164 Sydney, Australia, AUST R 206328
Product Registration Holder: Fresenius Medical Care Malaysia Sdn. Bhd.
2nd Floor, Axis Technology Centre, Lot 13, Jalan 51A/225, 46100 Petaling Jaya, Selangor. Malaysia.
Reg. No. : MAL-XXXXXXXXX
Jauh Daripada Kanak-kanak. **Controlled Medicine/ Ubat Terkawal**

KOREA

멀티빅2칼륨액5 L
전문의약품

[원료약품 및 분량]

	성 분 명	분 량
A액	염화칼슘이수화물(EP)	4.410 g
	염화마그네슘육수화물(EP)	2.033 g
	염화칼륨(EP)	2.982 g
	포도당수화물 (EP) (무수포도당으로서)	22.00 g (20.00 g)
B액	염화나트륨(EP)	6.463 g
	탄산수소나트륨(EP)	3.104 g
혼합액	나트륨	140.0 mmol
	칼륨	2.0 mmol
	칼슘	1.50 mmol
	마그네슘	0.50 mmol
	염소	111.0 mmol
	탄산수소염	35.0 mmol
	포도당	5.55 mmol

[성상] 투명한 플라스틱백에 든 무색 투명한 액상 제제
[효능효과]
- 급성 신부전 환자의 연속 혈액여과 및 혈액투석여과 시 체액대용액으로 사용
- 급성 신부전 환자의 연속 혈액투석 시 투석액으로 사용
  - 이 액은 특히 고칼륨혈증의 가능성이 있는 환자에게 적응된다.

[용법·용량] 환자의 체액균형상태, 특적적는 체액균형상태 및 환자 혈액 중 여과량에 따라 투여량 결정. 기타 상세 용법용량은 첨부문서 참조

[사용상의 주의사항]
- 다음 환자에는 투여하지 말 것.
  - 대사성 알칼리증 환자 - 저칼륨혈증 환자
- 기타 상세 사용상의 주의사항은 첨부문서 참조

[저장방법] 밀봉중기, 4~30℃에서 보관
제조자: Fresenius Medical Care Deutschland GmbH
Frankfurter Str. 6-8, 66606 St. Wendel, Germany
수입자: (주)프레제니우스에디컬케어코리아, 서울시 강남구 강남대로 308 랜드마크타워 7층 ☎: 02-2112-8878

multiBic®
2 mmol/L potassium

Na^+	140	mmol/L
K^+	**2.0**	**mmol/L**
Ca^{++}	1.5	mmol/L
Mg^{++}	0.50	mmol/L
Cl^-	111	mmol/L
HCO_3^-	35	mmol/L
Glucose (1g/L)	5.55	mmol/L
pH = 7.2		
Theoretical Osmolarity	296	mosmol/L

Isotonic solution

5000 mL X 2

Date of Manufacturing
사용기한/Expiry date
제조번호/Batch No.

69606080

multiBic® 2 mmol/L potassium
Marketing Authorisation Holder : **Fresenius Medical Care Deutschland GmbH, 61346 Bad Homburg v.d.H., Germany**
Manufactured by: **Fresenius Medical Care Deutschland GmbH, Frankfurter Str. 6–8, 66606 St. Wendel, Germany**

5000 mL x 2 Ref F00001151

FRESENIUS MEDICAL CARE

종이

Pantone 293C

211mm
34mm
25mm
5