



●SWKD

○WKMS

○WKMM

○HU

○HU (KK)

File Name: Ai Kan Shan_Leaflet-021220

Received : 15.11.2018

Amended : 02.12.2020

Output : 02.12.2020

Request : Shuying

Designer : Syafiq

Opened : 140w x 200h MM

Closed : 70w x 50h MM

Finished: -

Colour : 1C

K



Three Legs Ai Kan Shan Fever Powder

Paracetamol 500mg/sachet

Ubat Serbuk Demam Cap Kaki Tiga

Paracetamol 500mg/bungkus

三脚标外感散扑热息痛500毫克/包

Product Name:

Three Legs Ai Kan Shan Fever Powder

Paracetamol 500mg/sachet

Composition:

Paracetamol 500mg

Description:

White, fine powder. Odourless or almost odourless with a slight bitter taste.

Indication:

For relief of fever, headache, toothache and mild to moderate pain.

Dosage:

Adult: 1- 2 sachet every 4 hours. Maximum 8 sachets a day. To be taken after meal. Dissolve fever powder in a glass of water before taking.

Route of Administration:

Oral

Contraindication:

Nephropathy. Hypersensitivity to paracetamol.

Warning and Precaution:

This preparation contains paracetamol. Do not take any other paracetamol containing medicines at the same time. Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be sign of a serious condition. If these reactions occur, stop use and seek medical assistance right away. Paracetamol should be given with care to patients with impaired kidney or liver functions. Chronic use should be avoided. Daily use may potentiate oral anticoagulants.

Drug Interaction:

The speed of absorption of paracetamol maybe increased by metoclopramide or domperidone and absorption reduced by colestyramine. The anticoagulant effect of warfarin and other coumarins may be enhanced by prolonged regular use of paracetamol with increased risk of bleeding; occasional doses have no significant effect.

Pregnancy and Lactation:

Use in pregnancy:

- Considered to be the analgesic of choice in pregnant patients.
- Although it crosses placenta, paracetamol is considered to be safe in normal therapeutic doses for short-term use as a minor analgesic/antipyretic in pregnancy.

Use in lactation:

- Excreted in breast milk.
- Maternal ingestion of paracetamol in normal therapeutic doses does not appear to present a risk to the nursing infant.

Side Effects:

Side effects of paracetamol are usually mild, skin rashes and allergic reactions occur occasionally. Cutaneous hypersensitivity reactions including skin rashes, angioedema, Steven Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

Symptoms and Treatment of Overdose:

Symptoms : Nausea, vomiting, anorexia and abdominal pain, acute renal failure may develop. In severe poisoning, hepatic failure may proceed to encephalopathy, coma and death. Treatment : Remove the ingested drug by activated charcoal orally or N-acetylcysteine 150mg per kg body weight as a 20% solution given intravenously over 15 minutes followed by I.V. infusion of 50mg per kg in 500ml of dextrose injection over the next 4 hours and then 100mg per kg in one litre of dextrose injection over the next 16 hours.

Pharmacodynamics:

Paracetamol has an analgesic and antipyretic effects. It alleviates pain by inhibiting the prostaglandin synthesis of the brain, Therefore block the synthesis of prostaglandin. However, it will not alleviate any pain that arises from smooth muscle spasm in hollow viscera. It reduces fever by inhibiting the action of endogenous pyrogen on the hypothalamic heat-regulating centers to increase dissipation of body heat.

Pharmacokinetics:

Paracetamol is rapidly and completely absorbed from the gastro-intestinal system. The plasma concentration reaches a peak in ½ to 1 hour, and the plasma half-life is about 1 to 4 hours. Paracetamol is metabolized in the liver. The hydroxylated metabolites are responsible for methemoglobin formation and hepatotoxicity. About 3% of the absorbed paracetamol is excreted unchanged in the urine and about 80% in conjugated forms.

Storage Condition:

Store below 30°C. Protect from light and moisture. Keep out of reach of children.

Shelf Life:

5 years

Dosage Form:

Powder

Packing Size:

1.0g/sachet

The information contained in this leaflet is limited. For further information, please consult your pharmacist or doctor.

Nama Produk:

Ubat Serbuk Demam Cap Kaki Tiga

Paracetamol 500mg/bungkus

Komposisi:

Paracetamol 500mg

Deskripsi:

Serbuk halus yang berwarna putih, tidak berbau atau hampir tidak berbau dengan sedikit rasa pahit.

Indikasi:

Untuk melegakan demam, sakit kepala, sakit gigi dan kesakitan ringan.

Cara Penggunaan:

Dewasa: 1- 2 bungkus setiap 4 jam. Maksima 8 bungkus sehari. Ambil selepas makan. Larutkan ubat serbuk ini dalam segelas air dan minum.

Laluan Pemberian Ubat:

Oral

Kontraindikasi:

Pesakit buah pinggang dan yang hipersensitif terhadap paracetamol.

Amaran dan Perhatian:

Ubat in mengandungi paracetamol. Jangan makan ubat lain yang mengandungi paracetamol bersama dengan ubat ini. Alahan amaran: Paracetamol boleh menyebabkan alahan kulit yang teruk. Gejala boleh termasuk kemerahan kulit, lepuh atau ruam. Ini boleh menjadi tanda keadaan serius. Jika tindak balas ini berlaku, berhenti menggunakannya dan dapatkan bantuan perubatan dengan segera. Perlu diberikan dengan berhati-hati untuk pesakit yang mempunyai penjejasan pada buah pinggang atau hati. Penggunaan jangka panjang harus dielakkan. Penggunaan harian boleh meningkat kesan antikoagulan.

Interaksi dengan ubat lain:

Sila rujuk kepada penerangan Bahasa Inggeris.

Kehamilan dan Penyusuan:

Sila rujuk kepada penerangan Bahasa Inggeris.

Kesan Sampingan:

Sila rujuk kepada penerangan Bahasa Inggeris.

Symptom dan Rawatan untuk Kelebihan Dos:

Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakodinamik:

Sila rujuk kepada penerangan Bahasa Inggeris.

Farmakokinetik:

Sila rujuk kepada penerangan Bahasa Inggeris.

Kedaaan Penyimpanan:

Simpan di bawah 30°C. Lindungi daripada cahaya dan kelembapan. Jauhkan daripada capaian kanak-kanak.

Tempoh Hayat:

5 tahun.

Dos:

Serbuk

Saiz Bungkus:

1.0g / Bungkus

Maklumat yang terkandung dalam risalah ini adalah terhad. Untuk maklumat yang terperinci, sila hubungi ahli farmasi atau doktor.

产品名称:

三脚标外感散扑热息痛500毫克/包

成份:

扑热息痛500毫克

外观:

带有些许苦味. 无臭或近乎无臭的白色粉末。

主治:

解除发烧、头痛、牙痛及轻微疼痛。

剂量:

成人：每隔4小时1至2包。每日最多可服用8包。饭后服用。将药粉溶解在1杯开水后服食。

给药途径:

口服。

禁忌与注意:

肾病患者及对扑热息痛过敏者。

注意与防范:

此药含有扑热息痛，不可与别种含有扑热息痛的药一起服用。

过敏警示:

扑热息痛可能会引起严重的皮肤反应。症状包括皮肤发红，水泡或皮疹。这可能是病情严重的迹象。如果出现这些反应，请停止使用，并寻求医疗援助。扑热息痛应谨慎给予肾功能或肝功能受损的患者。应避免长期使用。日常使用中，可能会增加口服抗凝血药物。

药物相互作用:

请参阅英文版本。

怀孕和哺乳:

请参阅英文版本。

副作用:

请参阅英文版本。

药物过量症状及治疗:

请参阅英文版本。

药效学:

请参阅英文版本。

药代动力学:

请参阅英文版本。

贮存:

储藏低于30°C。避免阳光直射及潮湿处。远离儿童。

有效期:

5年

剂型:

粉末

包装:

1.0克/包

此说明书仅属略述，欲知更详细资料，可咨询您的药剂师或医生。

Manufactured by Product Registration Holder:

Syarikat Wen Ken Drug Sdn Bhd

24, Jalan Lambak, Taman Johor, 81200 Johor Bahru, Johor, Malaysia.

Singapore Distributor:

Wen Ken Marketing (S) Pte Ltd

2, Alexandra Road, #02-08, Delta House, Singapore 159919.

Date of Revision:

02.12.2020

...