

AXIMOL 650MG CAPLET

Paracetamol 650mg

What is in this leaflet

1. What Aximol is used for
2. How Aximol works
3. Before you use Aximol
4. How to use Aximol
5. While you are using it
6. Side effects
7. Storage and Disposal of Aximol
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What Aximol is used for

It is used for relief of headache, fever and mild to moderate pain.

How Aximol works

It contains paracetamol which has an analgesic and antipyretic effects that can alleviate pain by inhibiting the prostaglandin synthesis of the brain.

Before you use Aximol

- When you must not use it

Do not use it if you are allergic to paracetamol or other ingredients of Aximol.

Do not take any other paracetamol containing medicines at the same time.

- Before you start to use it

Safety for use in pregnancy and during breastfeeding has not been established. Inform your doctor if you are pregnant, planning to become pregnant or breastfeeding.

- Taking other medicines

Do not take any other paracetamol containing medicines at the same time.

Paracetamol should also be given with care to patients taking other drugs that affect the liver.

A combination of paracetamol and codeine has also enhanced warfarin anticoagulation activity.

Please inform the doctor or pharmacist if taking other medicines.

How to use Aximol

- How much to use

Adult: 1- ½ caplet every 4- 6 hours. Maximum 6 caplets a day.

6- 12 years old: ½ caplet every 4- 6 hours. Maximum 3 caplets a day.

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to use it

Please follow the advice of the doctor or pharmacist.

- How long to use it

Please check with your doctor or pharmacist.

- If you use too much (overdose)

If you have used too much Aximol, contact your doctor, an emergency department or a poisoning center immediately.

While you are using Aximol

- Things you must do

Inform your doctor or pharmacist of all the medications you are taking.

- Things you must not do

Do not take any new medicines without consulting your doctor.

Do not give Aximol to anyone else, even if they have the same symptoms or condition as you.

Side effects

Side-effects of paracetamol are usually mild, though haematological reactions such as agranulocytosis, thrombocytopenia and pancytopenia have been reported. Skin rashes and other allergic reactions occur occasionally. Cutaneous hypersensitivity reactions including skin rashes, angioedema, Stevens Johnson Syndrome/ Toxic Epidermal Necrolysis have been reported.

Allergy alert: Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters or rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.bpfk.gov.my (Consumers → Reporting).

Storage and Disposal of Aximol

- Storage

Store at room temperature below 30°C. Keep out of reach of children.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Oblong, convex, white, plain tablet with single score on one side only.

- Ingredients

- Active ingredient
Paracetamol
- Inactive ingredients
Isopropyl alcohol, corn starch, magnesium stearate, sodium starch glycolate, polyvinylpyrrolidone K90

- MAL number:

MAL10033158XCZ

Manufacturer

Sunward Pharmaceutical Sdn Bhd
No. 9, 11 & 17, Jalan Kempas 4, Taman Perindustrian Tampoi Indah, 81200 Johor Bahru, Johor, Malaysia.

Product Registration Holder

Syarikat Wen Ken Drug Sdn Bhd
24, Jalan Lambak, Taman Johor, 81200 Johor Bahru, Johor, Malaysia.

Date of revision

28/2/2017