THREE LEGS WOUND-KARE ANTISEPTIC CREAM

Cetrimide 0.5%w/w, Chlorhexidine Gluconate 0.1%w/w

What is in this leaflet

1. What Three Legs Wound-Kare Antiseptic Cream is used for
2. How Three Legs Wound-Kare Antiseptic Cream works
3. Before you used Three Legs Wound-Kare Antiseptic Cream
4. How to use Three Legs Wound-Kare Antiseptic Cream
5. While you are using it
6. Side effects
7. Storage and disposal of Three Legs Wound-Kare Antiseptic Cream
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial Number

What Three Legs Wound-Kare Antiseptic Cream is used for

Antiseptic and cleansing cream for use on cuts, scratches, blisters, grazes, windburn, sunburn, nappy rash, insect bites, cracked and itchy skin.

How Three Legs Wound-Kare Antiseptic Cream works

Chlorhexidine gluconate is an effective antiseptic with a wide range of activity against micro-organisms, including gram-positive and gram-negative bacteria, fungi, and viruses.
Cetrimide is a quaternary ammonium compound with surfactant and antiseptic properties.

Before you use Three Legs Wound-Kare Antiseptic Cream

- *When you must not use it*

Do not use it if you are allergic to any ingredients of Three Legs Wound-Kare Antiseptic Cream.

- *Before you start to use it*

Three Legs Wound-Kare Antiseptic Cream contains chlorhexidine.
Chlorhexidine is known to induce hypersensitivity, including generalized allergic reactions and anaphylactic shock.
Inform your healthcare provider if you have a known allergy to Chlorhexidine.
Stop using this product and seek immediate medical assistance if you experience rash, itching, swelling, breathing difficulties, light headedness or rapid heartbeat.
Cetrimide may have a prolonging effect on the wound healing process.

Three Legs Wound-Kare Antiseptic Cream is incompatible with anionic substances (e.g. soap).

Pregnancy and Lactation

There are no adequate data from the use of chlorhexidine gluconate and cetrimide in pregnant women.

It is not known whether chlorhexidine gluconate and cetrimide are excreted in breast milk. There is no adequate data from the use of chlorhexidine and cetrimide in breast-feeding women. However, it is unlikely that the products are excreted in breast milk since the products are poorly absorbed. After topical usage of the product, as a general precaution, rinse nipples thoroughly with water before breast-feeding.

- *Taking other medicines*

Please inform the doctor or pharmacist if taking other medicines.

How to use Three Legs Wound-Kare Antiseptic Cream

- *How much to use*

Apply gently onto the affected area 2-3 times daily. Wash hand thoroughly before and after each application.

For external use only. Not to be taken.

- *When to use it*

Please follow the advice of the doctor or pharmacist.

- *How long to use it*

Please follow the advice of the doctor or pharmacist.

- *If you forget to use it*

Continue to use it as directed by your doctor or on the package label.

- *If you use too much (overdose)*

While accidental ingestion is unlikely to cause any systemic effects due to poor absorption of chlorhexidine gluconate and cetrimide, ingestion of high concentrations may cause esophageal damage and necrosis with symptoms such as nausea and vomiting.

Treatment of poisoning is symptomatic; demulcents and diluents may be given if necessary, but emesis and lavage should be avoided.
Activated charcoal may be considered if the patient presents within an hour of ingestion. Corticosteroids may reduce oropharyngeal edema.

While you are using it

- *Things you must do*

Inform your doctor or pharmacist of all the medications you are taking.
Avoid contact with the eyes.

- *Things you must not do*

Do not take any new medicines without consulting your doctor or pharmacist.

- *Things to be careful of*

Ask your health care provider any question you may have about how to use Three Legs Wound-Kare Antiseptic Cream.

Side effects

Irritative skin reactions and hypersensitivity reactions to chlorhexidine have been reported. In the event that these reactions occur, discontinue use.
Chlorhexidine may cause anaphylaxis.
Cetrimide may have a prolonging effect on the wound healing process.
Immune system disorders
Frequently not known: Hypersensitivity including anaphylactic shock.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine. You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my (Consumer → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI).

THREE LEGS WOUND-KARE ANTISEPTIC CREAM

Cetrimide 0.5%w/w, Chlorhexidine Gluconate 0.1%w/w

Storage and Disposal of Three Legs Wound-Kare Antiseptic Cream

- *Storage*

Store below 30°C. Protect from direct sunlight. Keep out of reach of children.

- *Disposal*

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- *What it looks like*

White colour, smooth application and sticky cream with characteristic odour.

- *Ingredients*
- Active ingredients

Cetrimide, Chlorhexidine Gluconate

- Inactive ingredients

Chlorocresol, Cetostearyl Alcohol, Liquid Paraffin, White Soft Paraffin, Cetomacrogol 1000, Propylene Glycol & Purified Water

- *MAL number*

MALXXXXXX

Manufactured by Product Registration Holder

Syarikat Wen Ken Drug Sdn Bhd
24, Jalan Lambak,
Taman Johor,
81200 Johor Bahru,
Johor, Malaysia.

Date of revision

24/06/2024

Serial Number:

NPRA(R2)23/250