

DUORIDE-T 0.5MG/0.4MG CAPSULES

Dutasteride/Tamsulosin Hydrochloride (0.5mg/0.4mg)

What is in this leaflet

1. What Duoride-T is used for
2. How Duoride-T works
3. Before you use Duoride-T
4. How to use Duoride-T
5. While you are using Duoride-T
6. Side effects
7. Storage and Disposal of Duoride-T
8. Product description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial number

What Duoride-T is used for

Duoride-T is used to treat men with moderate to severe symptoms of benign prostatic hyperplasia (BPH) - a non-cancerous growth of the prostate gland caused by producing too much of a hormone called DHT (dihydrotestosterone). Duoride-T reduces the risk of acute urinary retention and the need for surgery in people with moderate to severe symptoms of BPH.

How Duoride-T works

Duoride-T capsules are a combination of two different medicines called dutasteride and tamsulosin. Dutasteride belongs to a group of medicines called 5-alpha reductase inhibitors, and tamsulosin belongs to a group of medicines called alpha-blockers.

As the prostate grows, it can lead to urinary problems, such as difficulty in passing urine and a need to go to the toilet more frequently. It can also cause the flow of the urine to be slower and less forceful. If left untreated, there is a risk that your urine flow will be completely blocked.

Dutasteride helps to shrink the prostate and relieve symptoms, by reducing the amount of DHT your body produces.

Tamsulosin acts by relaxing the muscles in your prostate gland, making it easier to pass urine, and improving your symptoms.

Before you use Duoride-T

– When you must not use it

Do not take Duoride-T:

- if you are allergic (hypersensitive) to dutasteride, tamsulosin or to any of the other ingredients of Duoride-T, or to other medicines known as 5-alpha reductase inhibitors.
- if you are female or age 21 years and below. This medicine is for adult men only.

If any of these apply to you, don't take Duoride-T until you have checked with your doctor.

Duoride-T must also not be taken by individuals with low blood pressure associated with feelings of dizziness upon getting up or standing (orthostatic hypotension), or by individuals with severe liver impairment.

– Before you start to use it

Take special care with Duoride-T

- Some individuals taking dutasteride and an alpha-blocker had heart failure more often than individuals taking only dutasteride or only an alpha-blocker. (Heart failure means your heart does not pump blood as well as it should.) Caution is advised in individuals with coronary heart disease and swallowing disorder.
- If you have mild to moderate liver disease, check with your doctor that Duoride-T is suitable for you. You may need extra check-ups while you are taking Duoride-T.
- If you have severe kidney impairment, incomplete bladder emptying and/or severely impaired urine flow.
- If you are going to have cataract (cloudy lens), glaucoma (increased eye pressure) surgery. Tell your doctor or eye specialist before your operation that you are taking, or have previously taken Duoride-T. They may ask you to temporarily stop taking Duoride-T 1-2 weeks before your operation.
- Women, children and adolescents must not handle leaking Duoride-T capsules, because the active ingredient can be absorbed through the skin. Wash the affected area immediately with soap and water if

there is any contact with the skin.

- Duoride-T may cause breast enlargement and tenderness. If this becomes troublesome, or if you notice breast lumps or nipple discharge you should talk to your doctor about these changes as these may be signs of a serious condition, such as breast cancer.

– Taking other medicines

Tell your doctor if you're taking any other medicines, or have recently taken any. This includes medicines bought without a prescription.

Don't take Duoride-T with these medicines:

- other alpha blockers (for enlarged prostate or high blood pressure)

Duoride-T is not recommended with these medicines:

- ketoconazole, itraconazole, fluconazole (used to treat fungal infections)

Some medicines can interact with Duoride-T and may make it more likely that you'll have side-effects. These medicines include:

- PDE5 inhibitors (used to help achieve or maintain an erection).
- warfarin (to prevent blood clotting)
- erythromycin, clarithromycin (antibiotics used to treat infections)
- verapamil, diltiazem, amiodarone (to treat irregular heartbeat)
- cimetidine (to treat heartburn)
- paroxetine (an antidepressant)

This is not a complete list of interactions. Tell your doctor or pharmacist if you are taking any medicines.

How to use Duoride-T

Always take Duoride-T exactly as your doctor has told you. Check with your doctor or pharmacist if you're not sure.

– How much to use

The usual dose of Duoride-T is one capsule (containing 0.5 mg dutasteride/0.4 mg tamsulosin) taken once a day.

DUORIDE-T 0.5MG/0.4MG CAPSULES

Dutasteride/Tamsulosin Hydrochloride (0.5mg/0.4mg)

– When to use it

Swallow your Duoride-T capsules whole with some water, approximately 30 minutes after the same meal each day. Don't chew or open the capsules. Contact with the contents of the capsules may make your mouth or throat sore.

– How long to use it

Take Duoride-T for as long as your doctor recommends. Don't stop taking Duoride-T without talking to your doctor first.

On discontinuing Duoride-T the prostate can go back to the size it was prior to the treatment.

– If you forget to use it

Don't take extra capsules to make up for a missed dose. Just take your next dose at the usual time.

– If you use too much (overdose)

If you take too many Duoride-T capsules, contact your doctor or pharmacist for advice. If possible, show them the Duoride-T pack. Acute low blood pressure (systolic blood pressure 70 mmHg), vomiting and diarrhea were observed in these people.

While you are using it

– Things you must do

Men taking dutasteride had a serious form of prostate cancer more often than men who did not take dutasteride.

A blood test to measure the amount of a substance called PSA (prostate specific antigen) in your blood can help your doctor to tell if you have prostate disease, including prostate cancer. If you have a higher than normal amount of PSA in your blood it could mean that you have prostate cancer.

Men taking Duoride-T should have their PSA measured 6 months after starting treatment and regularly thereafter.

Taking Duoride-T will reduce the amount of PSA measured in your blood. You could still be at risk for prostate cancer even though your PSA is lower. Your doctor can still use PSA to help detect prostate cancer, by comparing your test results each time you have a PSA test.

It is important to take your medicine as your doctor recommends. If you do not take it regularly it may interfere with your doctor's ability to monitor your PSA.

– Things you must not do

This medicine has been prescribed for you personally. Don't pass it on to other people it may harm them even if their symptoms seem to be the same as yours.

– Things to be careful of *Pregnancy and breast-feeding*

Women who are pregnant (or may be) must not handle leaking capsules. Dutasteride is absorbed through the skin and can affect the normal development of a male baby. Contact your doctor for advice if a pregnant woman has come into contact with the contents of a Duoride-T capsule.

Male Fertility

Duoride-T has been shown to reduce sperm count, semen volume and sperm movement. However, it is not clear if male fertility is affected by Duoride-T.

Driving and using machines

Duoride-T can make some people feel dizzy and have vision disturbance.

Don't drive or operate machinery unless you're feeling well.

Side Effects

Like all medicines, Duoride-T can cause side effects, but not everybody gets them.

Dizziness, light-headedness and fainting: Duoride-T can cause dizziness, light-headedness and on rare occasions fainting. Take care when moving from a lying down or sitting position to sitting or standing, particularly if you wake up in the night, until you know how

this medicine affects you. If you feel dizzy or light-headed at any time during treatment, sit or lie down until the symptoms pass.

Allergic reactions: Signs can include:

- raised and itchy rash(hives)
- swelling, sometimes of the face or mouth (angioedema) causing difficulty in breathing
- collapse.

Contact a doctor immediately if you get any of these symptoms. Stop taking Duoride-T.

Serious skin reactions: Signs can include:

- a widespread rash with blisters and peeling skin, particularly around the mouth, nose, eyes and genitals (Stevens-Johnson syndrome).

Contact a doctor immediately if you get these symptoms. Stop taking Duoride-T.

Common side effects:

- not able to achieve or maintain an erection
- decreased sex drive*
- difficulty with ejaculation*
- breast enlargement or tenderness

* these events may continue after you stop taking Duoride-T.

Uncommon side effects:

- Feeling of fast heart beat
- constipation, diarrhoea, vomiting, nausea
- weakness or loss of strength
- itchy, blocked or runny nose

Very rare side effects:

- persistent painful erection of the penis

Contact a doctor urgently if this happens to you. You may need to have treatment to avoid serious complications.

- depressed mood
- pain and swelling in your testicles.

Other side effects Other side effects have occurred but their exact frequency is unknown:

- irregular or fast heartbeat
- shortness of breath

DUORIDE-T 0.5MG/0.4MG CAPSULES

Dutasteride/Tamsulosin Hydrochloride (0.5mg/0.4mg)

- breast cancer in men

If you get side effects

Tell your doctor or pharmacist if any of the side effects listed becomes severe or troublesome, or if you notice any side effects not listed in this leaflet.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)]

Storage and Disposal of Duoride-T

– Storage

Keep out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and the label after EXP. The expiry date refers to the last day of that month.

Store below 30°C.

– Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

– What it looks like

This medicine is an oblong hard capsule (of around 24mm length) with a brown body and an orange cap.

Each hard capsules contains one dutasteride soft gelatin capsule and tamsulosin hydrochloride modified-pellets.

– Ingredients

– Active ingredient

The active substances of Duoride-T are Dutasteride 0.5mg and Tamsulosin hydrochloride 0.4mg.

– Inactive ingredient

- Glycerol monocaprylocaprate

- Butylhydroxytoluene
- Gelatin
- Glycerol
- Titanium Dioxide
- Yellow Iron Oxide
- Triglycerides
- Lecithin (soya)
- Microcrystalline Cellulose
- Magnesium Stearate
- Methacrylic acid-Ethyl acrylate copolymer
- Sodium Hydroxide
- Triacetin
- Purified Talc
- Titanium Dioxide
- Talc

– MAL numbers

MALXXXXXXXXXX

Manufacturers

SAG MANUFACTURING,
S.L.U.

Carretera Nacional 1 Km 36,
San Agustín del Guadalix 28750
Madrid – Spain

Product Registration Holder

Novugen Pharma Sdn. Bhd.
No. 3, Jalan Jururancang U1/21,
Hicom Glenmarie Industrial Park,
40150, Shah Alam,
Selangor, Malaysia.

Date of revision

26/10/2022

Serial Number

NPRA (R2) 22/091