




50mm

47mm

PEM-I/S2/P2-001/L3/M1-02

Exp.
Lot.
Mfg.

80736B03

ONTREX 500mg
Powder for Concentrate
for Solution for Infusion

Pemetrexed 500mg

ONTREX 500mg
Powder for Concentrate
for Solution for Infusion

Each vial contains pemetrexed disodium 2.5 hydrate equivalent to 500 mg pemetrexed, 500 mg of mannitol. Hydrochloric acid and/or sodium hydroxide may have been added to adjust pH.

To Reconstitute: Add 20 mL of 0.9% Sodium Chloride Injection (preservative free) to make a solution containing 25 mg/mL pemetrexed. Reconstituted solution must be further diluted before use (see prescribing information).

Administer infusion solution within 24 hours after initial reconstitution. Discard unused portion.

See prescribing information for dosage.
MAL18126003AZ

ONTREX 500mg
Powder for Concentrate
for Solution for Infusion

Pemetrexed 500mg

KEEP OUT OF REACH OF CHILDREN
JAUHI UBAT DARIPADA KANAK-KANAK



Storage: ONTREX 500mg powder for concentrate for solution for infusion should be stored below 30°C (86°F). Chemical and physical stability of reconstituted and infusion solutions of ONTREX 500mg powder for concentrate for solution for infusion were demonstrated for up to 24 hours following initial reconstitution, when stored refrigerated, 2° to 8°C (36° to 46°F). When prepared as directed, reconstituted and infusion solutions of ONTREX 500mg powder for concentrate for solution for infusion contain no antimicrobial preservatives. Discard unused portion.

CAUTION: CYTOTOXIC AGENT

Controlled Medicine

Manufactured by:
Jiangsu Hansoh Pharmaceutical Group Co., Ltd.,
Lianyungang, Jiangsu 222069, China.



Product Registration Holder:
Novugen Oncology Sdn. Bhd.
No. 3, Jalan Jururancang U1/21,
Hicom Glenmarie Industrial Park,
40150 Shah Alam, Selangor, Malaysia.

One Single-Use Vial (1 Vial)
For intravenous use only

80mm