Size : 35*35*65mm

IRI-I/S1/P2-001/L3/M1-07

Inotecam

(20mg/ml) Concentrate for Solution for Infusion

40mg/2ml

Irinotecan

For Intravenous Infusion Use Only

1 x 2mL Vial

(Single Dose)

Each vial contains irinotecan (as hydrochloride trihydrate), 40mg

See prescribing information for dosage.

Must be diluted before use.

KEEP MEDICINE OUT OF REACH OF CHILDREN
JAUHI UBAT DARIPADA KANAK-KANAK

CAUTION: CYTOTOXIC AGENT

Controlled Medicine

MAL XXXXXXXXXX

Inotecam

(20mg/ml) Concentrate for Solution for Infusion

40mg/2ml

Irinotecan

For Intravenous Infusion Use Only

1 x 2mL Vial

(Single Dose)

Storage: Store below 30°C. Protect from light and freezing. Retain in carton. Discard unused portion.

In-use storage condition: Solutions (0.12mg/mL, 2.8mg/mL) diluted in 5% Dextrose Injection is physically and chemically stable at room temperature (approximately 25°C) and in ambient fluorescent lighting within 24h, at refrigerated temperatures (2°C to 8°C) and protected from light within 48h. Solutions (0.12mg/mL, 2.8mg/mL) diluted in 0.9 % Sodium Chloride Injection are physically and chemically stable within 24 hours at room temperature (approximately 25°C) and in ambient fluorescent lighting.

IF CONTENTS ARE DAMAGE AND/OR LEAKING INCINERATE ENTIRE PACKAGE.

Manufactured by:
Qilu Pharmaceutical (Hainan) Co., Ltd.
Haikou, 570314, China.

Product Registration Holder:
Novugen Oncology Sdn. Bhd.
No.3, Jalan Jururancang U1/21,
Hicom Glenmarie Industrial Park,
40150 Shah Alam, Selangor, Malaysia.

Mfg.
Lot.
Exp.

Inotecam

(20mg/ml) Concentrate for Solution for Infusion

40mg/2ml

Irinotecan

For Intravenous Infusion Use Only

1 x 2mL Vial

(Single Dose)