


Unvarnished area for pasting

**A Step-by-Step Guide to taking your Cladribine Tablets 10 mg**

Cladribine tablets are packed in child-resistant blister packs inside a carton and must be kept out of the sight and reach of children. See below for a step-by-step guide on how to handle the package and to take the Cladribine tablets. Make sure you know how many tablets are contained in the package. See package leaflet for guidance.



**1** Have a glass of water ready and make sure your hands are clean and dry before taking the tablet(s).




**2** Open flap on carton.



**4** Peel back the label of the blister pack(s).



**6** Swallow tablet(s) with water. Tablets must be swallowed whole and not chewed or allowed to dissolve in your mouth. Contact with skin should be limited. Avoid touching your nose, eyes, and other parts of the body.



**7** Wash your hands thoroughly with soap and water.

DAY 1	DAY 2	DAY 3	DAY 4
1 tablet + 0	1 tablet + 0	1 tablet + 0	1 tablet + 0

Varnish free area for overcoding

**IMPORTANT: Contact Doctor if incorrect dose taken. Call your doctor for medical advice about side effects. You may report Side effects to FDA at 1-800-FDA-1088.**

**STORAGE and DISPOSAL:** Store at controlled room temperature, 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Store in original package in order to protect from moisture. Contact your doctor for instructions on proper disposal of unused Cladribine tablets. Keep out of reach of children.

Manufactured by:  
Novugen Oncology Sdn. Bhd.  
No 47 Jalan Lengkuk Teknologi 2,  
Taman Teknologi Enstek Fasa 1,  
Techpark@Enstek,  
71760 Labu,  
Negeri Sembilan Darul Khusus.

Manufactured for:  
Accord Healthcare, Inc.,  
Raleigh, NC 27617.



NDC 16729-625-02

**Accord Cladribine Tablets**

**10 mg per tablet**

Rx only

**Contents:**

4 tablets: Four child-resistant blister packs each containing one 10 mg tablet.

**Cytotoxic Agent:**

Handle tablets as directed.

Dispense with enclosed Medication Guide.

4 Tablets



NDC 16729-625-02  
**Accord Cladribine Tablets**  
**10 mg per tablet**  
Rx only  
Contents:  
4 tablets: Four child-resistant blister packs each containing one 10 mg tablet.  
**Cytotoxic Agent:**  
Handle tablets as directed.  
Dispense with enclosed Medication Guide.  
4 Tablets

Each Cladribine 10 mg tablet contains cladribine USP, hydroxypropyl betadex complex, sorbitol, and magnesium stearate. Cladribine Tablets are provided as 10 mg white to off white, round, biconvex uncoated tablets debossed with "TU" on one side and plain on the other side.

**Dosage and Administration:**

Based on body weight. For oral administration.

See enclosed Prescribing Information and Medication Guide.

**Separate ingestion of other oral medicines by at least 3 hours before and after taking Cladribine Tablets.**

**SPECIAL HANDLING:** Remove tablet from foil blister just prior to use. Immediately swallow tablet whole with water. Do NOT chew. Avoid prolonged contact with skin. Wash hands and surfaces.

The unit dose blister pack is child-resistant.

File Name : Cladribine-10mg(Novugen)4T-Box

Size : 80 x 40 x 56 (mm)

Colour : Pantone Orange 021 C, 166 C, 430 C + CMYK

Date : 01/08/25 (1), 05/08/25, 12/01/26


Unvarnished area for pasting

A Step-by-Step Guide to taking your Cladribine Tablets 10 mg

Cladribine tablets are packed in child-resistant blister packs inside a carton and must be kept out of the sight and reach of children. See below for a step-by-step guide on how to handle the package and to take the Cladribine tablets. Make sure you know how many tablets are contained in the package. See package leaflet for guidance.



1 Have a glass of water ready and make sure your hands are clean and dry before taking the tablet(s).



2 Open flap on carton.



4 Peel back the label of the blister pack(s).



6 Swallow tablet(s) with water. Tablets must be swallowed whole and not chewed or allowed to dissolve in your mouth. Contact with skin should be limited. Avoid touching your nose, eyes, and other parts of the body.



7 Wash your hands thoroughly with soap and water.

DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
1 tablet	1 tablet	1 tablet	1 tablet	1 tablet

Varnish free area for overcoding

**IMPORTANT: Contact Doctor if incorrect dose taken. Call your doctor for medical advice about side effects. You may report Side effects to FDA at 1-800-FDA-1088.**

**STORAGE and DISPOSAL:** Store at controlled room temperature, 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Store in original package in order to protect from moisture. Contact your doctor for instructions on proper disposal of unused Cladribine tablets. Keep out of reach of children.

Manufactured by:  
Novugen Oncology Sdn. Bhd.  
No 47 Jalan Lengkok Teknologi 2,  
Taman Teknologi Enstek Fasa 1,  
Techpark@Enstek,  
71760 Labu,  
Negeri Sembilan Darul Khusus.

Manufactured for:  
Accord Healthcare, Inc.,  
Raleigh, NC 27617.



NDC 16729-625-32

Accord Cladribine Tablets

10 mg per tablet

Rx only

Contents:  
5 tablets: Five child-resistant blister packs each containing one 10 mg tablet.

**Cytotoxic Agent:**

Handle tablets as directed.

Dispense with enclosed Medication Guide.

5 Tablets



Each Cladribine 10 mg tablet contains cladribine USP, hydroxypropyl betadex complex, sorbitol, and magnesium stearate. Cladribine Tablets are provided as 10 mg white to off white, round, biconvex uncoated tablets debossed with "TU" on one side and plain on the other side.

**Dosage and Administration:**

Based on body weight. For oral administration. See enclosed Prescribing Information and Medication Guide.

**Separate ingestion of other oral medicines by at least 3 hours before and after taking Cladribine Tablets.**

**SPECIAL HANDLING:** Remove tablet from foil blister just prior to use. Immediately swallow tablet whole with water. Do NOT chew. Avoid prolonged contact with skin. Wash hands and surfaces.

The unit dose blister pack is child-resistant.

Rx only  
Contents:  
5 tablets: Five child-resistant blister packs each containing one 10 mg tablet.  
**Cytotoxic Agent:**  
Handle tablets as directed.  
Dispense with enclosed Medication Guide.  
5 Tablets

File Name : Cladribine-10mg(Novugen)5T-Box

Size : 80 x 46 x 56 (mm)

Colour : Pantone 7476 C, 166 C, 430 C + CMYK

Date : 01/08/25 (1), 05/08/25, 12/01/26

Through cuts length  
4 mm at a distance of  
7 mm + Creasing is required

11 mm

10 mm


### Unvarnished area for pasting

#### A Step-by-Step Guide to taking your Cladribine Tablets 10 mg

Cladribine tablets are packed in child-resistant blister packs inside a carton and must be kept out of the sight and reach of children. See below for a step-by-step guide on how to handle the package and to take the Cladribine tablets. Make sure you know how many tablets are contained in the package. See package leaflet for guidance.



1 Have a glass of water ready and make sure your hands are clean and dry before taking the tablet(s).



2 Open flap on carton.



4 Peel back the label of the blister pack(s).

3 Take out the number of blister packs needed for your prescribed dose.

5 Place 1 or 2 tablet(s) into your hand, according to your prescribed dose.



6 Swallow tablet(s) with water. Tablets must be swallowed whole and not chewed or allowed to dissolve in your mouth. Contact with skin should be limited. Avoid touching your nose, eyes, and other parts of the body.



7 Wash your hands thoroughly with soap and water.

DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
2 tablets	1 tablet	1 tablet	1 tablet	1 tablet

Varnish free area  
for overcoding

**IMPORTANT: Contact Doctor if incorrect dose taken. Call your doctor for medical advice about side effects. You may report Side effects to FDA at 1-800-FDA-1088.**

**STORAGE and DISPOSAL:** Store at controlled room temperature, 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Store in original package in order to protect from moisture. Contact your doctor for instructions on proper disposal of unused Cladribine tablets. Keep out of reach of children.

Manufactured by:  
Novugen Oncology Sdn. Bhd.  
No 47 Jalan Lengkok Teknologi 2,  
Taman Teknologi Enstek Fasa 1,  
Techpark@Enstek,  
71760 Labu,  
Negeri Sembilan Darul Khusus.



Manufactured for:  
Accord Healthcare, Inc.,  
Raleigh, NC 27617.

Through cuts length  
4 mm at a distance of  
7 mm + Creasing is required

**10 mg per tablet**  
Rx only  
Contents:  
6 tablets: Six child-resistant blister packs each containing one 10 mg tablet.  
**Cytotoxic Agent:**  
Handle tablets as directed.  
Dispense with enclosed Medication Guide.  
6 Tablets

**Accord Cladribine Tablets**

NDC 16729-625-66

NDC 16729-625-66

## Accord Cladribine Tablets

**10 mg per tablet**

Rx only

Contents:

6 tablets: Six child-resistant blister packs each containing one 10 mg tablet.

**Cytotoxic Agent:**

Handle tablets as directed.

Dispense with enclosed Medication Guide.

6 Tablets

accord

Through cuts length  
4 mm at a distance of  
7 mm + Creasing is required

Each Cladribine 10 mg tablet contains cladribine USP, hydroxypropyl betadex complex, sorbitol, and magnesium stearate. Cladribine Tablets are provided as 10 mg white to off white, round, biconvex uncoated tablets debossed with "TU" on one side and plain on the other side.

#### Dosage and Administration:

Based on body weight. For oral administration. See enclosed Prescribing Information and Medication Guide.

**Separate ingestion of other oral medicines by at least 3 hours before and after taking Cladribine Tablets.**

**SPECIAL HANDLING:** Remove tablet from foil blister just prior to use. Immediately swallow tablet whole with water. Do NOT chew. Avoid prolonged contact with skin. Wash hands and surfaces.

The unit dose blister pack is child-resistant.

Through cuts length  
4 mm at a distance of  
7 mm + Creasing is required


18 mm

File Name : Cladribine-10mg(Novugen)6T-Box

Size : 80 x 52 x 56 (mm)

Colour : Pantone 2577 C, 166 C, 430 C + CMYK

Date : 01/08/25 (1), 05/08/25, 12/01/26



Unvarnished area for pasting

**A Step-by-Step Guide to taking your Cladribine Tablets 10 mg**  
 Cladribine tablets are packed in child-resistant blister packs inside a carton and must be kept out of the sight and reach of children. See below for a step-by-step guide on how to handle the package and to take the Cladribine tablets. Make sure you know how many tablets are contained in the package. See package leaflet for guidance.

- 1 Have a glass of water ready and make sure your hands are clean and dry before taking the tablet(s).
- 2 Open flap on carton.
- 3 Take out the number of blister packs needed for your prescribed dose.
- 4 Peel back the label of the blister pack(s).
- 5 Place 1 or 2 tablet(s) into your hand, according to your prescribed dose.
- 6 Swallow tablet(s) with water. Tablets must be swallowed whole and not chewed or allowed to dissolve in your mouth. Contact with skin should be limited. Avoid touching your nose, eyes, and other parts of the body.
- 7 Wash your hands thoroughly with soap and water.

DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
2 tablets	2 tablets	1 tablet	1 tablet	1 tablet

**IMPORTANT: Contact Doctor if incorrect dose taken. Call your doctor for medical advice about side effects. You may report Side effects to FDA at 1-800-FDA-1088.**

Manufactured by:  
 Novugen Oncology Sdn. Bhd.  
 No 47, Jalan Lengkok Teknologi 2,  
 Taman Teknologi Enstek Fasa 1,  
 Techpark@Enstek,  
 71760 Labu,  
 Negeri Sembilan Darul Khusus.

**STORAGE and DISPOSAL:** Store at controlled room temperature, 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Store in original package in order to protect from moisture. Contact your doctor for instructions on proper disposal of unused Cladribine tablets. Keep out of reach of children.



Manufactured for:  
 Accord Healthcare, Inc.,  
 Raleigh, NC 27617.

Through cuts length 4 mm at a distance of 7 mm + Creasing is required

Varnish free area for overcoding

NDC 16729-625-86  
**Accord Cladribine Tablets**  
**10 mg per tablet**  
 Rx only  
 Contents:  
 7 tablets: Seven child-resistant blister packs each containing one 10 mg tablet.  
**Cytotoxic Agent:**  
 Handle tablets as directed.  
 Dispense with enclosed Medication Guide.  
 7 Tablets

NDC 16729-625-86  
**Accord Cladribine Tablets**

**10 mg per tablet**  
 Rx only  
 Contents:  
 7 tablets: Seven child-resistant blister packs each containing one 10 mg tablet.  
**Cytotoxic Agent:**  
 Handle tablets as directed.  
 Dispense with enclosed Medication Guide.  
 7 Tablets  
 accord

Through cuts length 4 mm at a distance of 7 mm + Creasing is required

Each Cladribine 10 mg tablet contains cladribine USP, hydroxypropyl betadex complex, sorbitol, and magnesium stearate. Cladribine Tablets are provided as 10 mg white to off white, round, biconvex uncoated tablets debossed with "TU" on one side and plain on the other side.

**Dosage and Administration:**  
 Based on body weight. For oral administration. See enclosed Prescribing Information and Medication Guide.  
**Separate ingestion of other oral medicines by at least 3 hours before and after taking Cladribine Tablets.**

**SPECIAL HANDLING:** Remove tablet from foil blister just prior to use. Immediately swallow tablet whole with water. Do NOT chew. Avoid prolonged contact with skin. Wash hands and surfaces.

The unit dose blister pack is child-resistant.

Through cuts length 4 mm at a distance of 7 mm + Creasing is required

11 mm

10 mm

Through cuts length 4 mm at a distance of 7 mm + Creasing is required

Unvarnished area for pasting

Varnish free area for overcoating

**A Step-by-Step Guide to taking your Cladribine Tablets 10 mg**

Cladribine tablets are packed in child-resistant blister packs inside a carton and must be kept out of the sight and reach of children. See below for a step-by-step guide on how to handle the package and to take the Cladribine tablets. Make sure you know how many tablets are contained in the package. See package leaflet for guidance.

- 1 Have a glass of water ready and make sure your hands are clean and dry before taking the tablet(s).
- 2 Open flap on carton.
- 3 Take out the number of blister packs needed for your prescribed dose.
- 4 Peel back the label of the blister pack(s).
- 5 Place 1 or 2 tablet(s) into your hand, according to your prescribed dose.
- 6 Swallow tablet(s) with water. Tablets must be swallowed whole and not chewed or allowed to dissolve in your mouth. Contact with skin should be limited. Avoid touching your nose, eyes, and other parts of the body.
- 7 Wash your hands thoroughly with soap and water.

DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
2 tablets	2 tablets	2 tablets	1 tablet	1 tablet

**IMPORTANT: Contact Doctor if incorrect dose taken. Call your doctor for medical advice about side effects. You may report Side effects to FDA at 1- 800-FDA-1088.**

**STORAGE and DISPOSAL:** Store at controlled room temperature, 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Store in original package in order to protect from moisture. Contact your doctor for instructions on proper disposal of unused Cladribine tablets. Keep out of reach of children.

Manufactured for:  
Accord Healthcare, Inc.,  
Raleigh, NC 27617.

Manufactured by:  
Novugen Oncology Sdn. Bhd.  
No 47 Jalan Lengkok Teknologi 2,  
Taman Teknologi Enstek Fasa 1,  
Techpark@Enstek,  
71760 Labu,  
Negeri Sembilan Darul Khusus.



Through cuts length 4 mm at a distance of 7 mm + Creasing is required

NDC 16729-625-65

**Accord Cladribine Tablets**

**10 mg per tablet**

Rx only

Contents: Eight child-resistant blister packs each containing one 10 mg tablet.

**Cytotoxic Agent:**

Handle tablets as directed.

Dispense with enclosed Medication Guide.

8 Tablets

NDC 16729-625-65

**Accord Cladribine Tablets**

**10 mg per tablet**

Rx only

Contents: 8 tablets: Eight child-resistant blister packs each containing one 10 mg tablet.

**Cytotoxic Agent:**

Handle tablets as directed.

Dispense with enclosed Medication Guide.

8 Tablets

Through cuts length 4 mm at a distance of 7 mm + Creasing is required

Each Cladribine 10 mg tablet contains cladribine USP, hydroxypropyl betadex complex, sorbitol, and magnesium stearate. Cladribine Tablets are provided as 10 mg white to off white, round, biconvex uncoated tablets debossed with "TU" on one side and plain on the other side.

**Dosage and Administration:**  
Based on body weight. For oral administration. See enclosed Prescribing Information and Medication Guide.  
**Separate ingestion of other oral medicines by at least 3 hours before and after taking Cladribine Tablets.**

**SPECIAL HANDLING:** Remove tablet from foil blister just prior to use. Immediately swallow tablet whole with water. Do NOT chew. Avoid prolonged contact with skin. Wash hands and surfaces.

The unit dose blister pack is child-resistant.

Through cuts length 4 mm at a distance of 7 mm + Creasing is required


20 mm

File Name : Cladribine-10mg(Novugen)8T-Box

Size : 80 x 64 x 56 (mm)

Colour : Pantone 1795 C, 166 C, 430 C + CMYK

Date : 01/08/25 (1), 05/08/25, 12/01/26



Unvarnished area for pasting

**A Step-by-Step Guide to taking your Cladribine Tablets 10 mg**  
 Cladribine tablets are packed in child-resistant blister packs inside a carton and must be kept out of the sight and reach of children. See below for a step-by-step guide on how to handle the package and to take the Cladribine tablets. Make sure you know how many tablets are contained in the package. See package leaflet for guidance.

- 1 Have a glass of water ready and make sure your hands are clean and dry before taking the tablet(s).
- 2 Open flap on carton.
- 3 Take out the number of blister packs needed for your prescribed dose.
- 4 Peel back the label of the blister pack(s).
- 5 Place 1 or 2 tablet(s) into your hand, according to your prescribed dose.
- 6 Swallow tablet(s) with water. Tablets must be swallowed whole and not chewed or allowed to dissolve in your mouth. Contact with skin should be limited. Avoid touching your nose, eyes, and other parts of the body.
- 7 Wash your hands thoroughly with soap and water.

DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
2 tablets	2 tablets	2 tablets	2 tablets	1 tablet

**IMPORTANT: Contact Doctor if incorrect dose taken.**  
**Call your doctor for medical advice about side effects.**  
**You may report Side effects to FDA at 1- 800-FDA-1088.**

**STORAGE and DISPOSAL:** Store at controlled room temperature, 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature].  
 Store in original package in order to protect from moisture. Contact your doctor for instructions on proper disposal of unused Cladribine tablets. Keep out of reach of children.

Manufactured for:  
 Accord Healthcare, Inc.,  
 Raleigh, NC 27617.

Manufactured by:  
 Novugen Oncology Sdn. Bhd.  
 No 47 Jalan Lengkuk Teknologi 2,  
 Taman Teknologi Enstek Fasa 1,  
 Techpark@Enstek,  
 71760 Labu,  
 Negeri Sembilan Darul Khusus.



Through cuts length  
 4 mm at a distance of  
 7 mm + Creasing is required

NDC 16729-625-34  
**Accord Cladribine Tablets**  
**10 mg per tablet**  
 Rx only  
**Contents:**  
 9 tablets: Nine child-resistant blister packs each containing one 10 mg tablet.  
**Cytotoxic Agent:**  
 Handle tablets as directed.  
 Dispense with enclosed Medication Guide.  
 9 Tablets

NDC 16729-625-34  
**Accord Cladribine Tablets**  
**10 mg per tablet**

Rx only  
**Contents:**  
 9 tablets: Nine child-resistant blister packs each containing one 10 mg tablet.  
**Cytotoxic Agent:**  
 Handle tablets as directed.  
 Dispense with enclosed Medication Guide.  
**9 Tablets**

Through cuts length  
 4 mm at a distance of  
 7 mm + Creasing is required

Each Cladribine 10 mg tablet contains cladribine USP, hydroxypropyl betadex complex, sorbitol, and magnesium stearate. Cladribine Tablets are provided as 10 mg white to off white, round, biconvex uncoated tablets debossed with "TU" on one side and plain on the other side.

**Dosage and Administration:**  
 Based on body weight. For oral administration. See enclosed Prescribing Information and Medication Guide.  
**Separate ingestion of other oral medicines by at least 3 hours before and after taking Cladribine Tablets.**

**SPECIAL HANDLING:** Remove tablet from foil blister just prior to use. Immediately swallow tablet whole with water. Do NOT chew. Avoid prolonged contact with skin. Wash hands and surfaces.

The unit dose blister pack is child-resistant.

Through cuts length  
 4 mm at a distance of  
 7 mm + Creasing is required

20 mm






Through cuts length  
4 mm at a distance of  
7 mm + Creasing is required

11 mm

10 mm

Unvarnished area for pasting

**A Step-by-Step Guide to taking your Cladribine Tablets 10 mg**  
Cladribine tablets are packed in child-resistant blister packs inside a carton and must be kept out of the sight and reach of children. See below for a step-by-step guide on how to handle the package and to take the Cladribine tablets. Make sure you know how many tablets are contained in the package. See package leaflet for guidance.

- 1 Have a glass of water ready and make sure your hands are clean and dry before taking the tablet(s).
- 2 Open flap on carton.
- 3 Take out the number of blister packs needed for your prescribed dose.
- 4 Peel back the label of the blister pack(s).
- 5 Place 1 or 2 tablet(s) into your hand, according to your prescribed dose.
- 6 Swallow tablet(s) with water. Tablets must be swallowed whole and not chewed or allowed to dissolve in your mouth. Contact with skin should be limited. Avoid touching your nose, eyes, and other parts of the body.
- 7 Wash your hands thoroughly with soap and water.

DAY 1	DAY 2	DAY 3	DAY 4	DAY 5
2 tablets	2 tablets	2 tablets	2 tablets	2 tablets

Varnish free area  
for overcoding

**IMPORTANT: Contact Doctor if incorrect dose taken.**  
**Call your doctor for medical advice about side effects.**  
**You may report Side effects to FDA at 1-800-FDA-1088.**

**STORAGE and DISPOSAL:** Store at controlled room temperature, 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [see USP Controlled Room Temperature]. Store in original package in order to protect from moisture. Contact your doctor for instructions on proper disposal of unused Cladribine tablets. Keep out of reach of children.

Manufactured for:  
Accord Healthcare, Inc.,  
Raleigh, NC 27617.

Manufactured by:  
Novugen Oncology Sdn. Bhd.  
No 47 Jalan Lengku Teknologi 2,  
Taman Teknologi Enstek Fasa 1,  
Techpark@Enstek,  
71760 Labu,  
Negeri Sembilan Darul Khusus.



Through cuts length  
4 mm at a distance of  
7 mm + Creasing is required

NDC 16729-625-03

## Accord Cladribine Tablets

**10 mg** per tablet

Rx only

**Contents:**  
10 tablets: Ten child-resistant blister packs each containing one 10 mg tablet.

**Cytotoxic Agent:**

**Handle tablets as directed.**

**Dispense with enclosed Medication Guide.**

10 Tablets

accord

Through cuts length  
4 mm at a distance of  
7 mm + Creasing is required

Each Cladribine 10 mg tablet contains cladribine USP, hydroxypropyl betadex complex, sorbitol, and magnesium stearate. Cladribine Tablets are provided as 10 mg white to off white, round, biconvex uncoated tablets debossed with "TU" on one side and plain on the other side.

**Dosage and Administration:**

Based on body weight. For oral administration.  
See enclosed Prescribing Information and Medication Guide.

**Separate ingestion of other oral medicines by at least 3 hours before and after taking Cladribine Tablets.**

**SPECIAL HANDLING:** Remove tablet from foil blister just prior to use. Immediately swallow tablet whole with water. Do NOT chew. Avoid prolonged contact with skin. Wash hands and surfaces.

The unit dose blister pack is child-resistant.

Through cuts length  
4 mm at a distance of  
7 mm + Creasing is required

20 mm

File Name : Cladribine-10mg(Novugen)10T-Box  
Size : 80 x 76 x 56 (mm)  
Colour : Pantone 266 C, 166 C, 430 C + CMYK  
Date : 01/08/25 (1), 05/08/25, 12/01/26