

GOFEN LIQUIZ SUSPENSION

Ibuprofen (100mg/5ml)

What is in this leaflet

1. What Gofen Liquiz is used for
2. How Gofen Liquiz works
3. Before you use Gofen Liquiz
4. How to use Gofen Liquiz
5. While you are using it
6. Side effects
7. Storage and Disposal of Gofen Liquiz
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of RiMUP revision

What Gofen Liquiz is used for

Gofen Liquiz is used to relief of fever associated with cold and flu, as well as pain from teething, toothache, earache, sore throats, headache, minor aches and sprains.

Also for the relief of inflammation of muscles and joints.

How Gofen Liquiz works

Gofen Liquiz contains ibuprofen, a non steroidal anti-inflammatory drug (NSAIDs). It works by reducing prostaglandins which is natural substance that causes pain and inflammation.

Before you use Gofen Liquiz

- When you must not use it

Do not take this medicine if:

- you are allergic to ibuprofen or any other ingredients of this medicine (listed in Product Description)
- you are hypersensitive to aspirin, ibuprofen or other nonsteroidal anti-inflammatory drugs.
- you have history of peptic ulceration.

- Before you start to use it

Check with your doctor or pharmacist before taking Gofen Liquiz if you have:

- asthma or allergies
- any gastrointestinal problem
- kidney, liver or heart problem
- high blood pressure
- developing gastrointestinal bleeding (heavy or frequent alcohol drinking, smoking or taking medications or

conditions that may increase chance of bleeding)

Pregnancy and lactation

Please consult your doctor or pharmacist if you are pregnant, planning for pregnancy or breast-feeding before using any medicine.

- Taking other medicines

Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or health food shop.

Inform your doctor or pharmacist if you are taking:

- Lithium
- Methotrexate
- Blood thinning medication
- High blood pressure medications
- Other pain relievers

How to use Gofen Liquiz

Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- How much to use

Infants 3-6months weighing more than 5kg: 2.5ml may be taken 3 times in 24 hours.

Infants 6-12months: 2.5ml may be taken 3 to 4 times in 24 hours.

Children 1-3 years: 5ml may be taken 3 times in 24 hours.

Children 4-6 years: 7.5ml may be taken 3 times in 24 hours.

Children 7-9 years: 10ml may be taken 3 times in 24 hours.

Children 10-12 years: 15ml may be taken 3 times in 24 hours.

- When to use it

Use as directed by your doctor or pharmacist.

- How long to use it

Use this medicine for the duration that your doctor recommends.

- If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you or anyone else may have use too much of this medicine. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

Taking too much of this medicine may cause headache, nausea, vomiting, drowsiness and loss of consciousness.

While you are using Gofen Liquiz

- Things you must do

Tell all the doctors, dentists and pharmacists treating you that you are using Gofen Liquiz.

- Things you must not do

Do not stop using the medicine unless advised by your doctor.

Do not use any new medicines without consulting your doctor or pharmacist.

Do not give Gofen Liquiz to anyone else, even if they have the same symptoms or condition as you.

- Things to be careful of

Driving and using machines

This medicine may affect your ability to drive or use machines. If this medicine makes you feel sick, dizzy or tired, do not drive or use machines and contact your doctor immediately.

Side effects

Like all medicines, Gofen Liquiz can cause side effects, although not everybody gets them.

Visit your doctor or pharmacist immediately if you experience any side effects after using this medicine.

Adverse reactions are usually mild and transient: Stomach discomfort, nausea, stomach ulcer, rashes or worsening of asthma, liver and kidney problems, dizziness, headache and hearing disturbance.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Gofen Liquiz

- Storage

Keep out of the reach and sight of children.

Store below 30 °C in a dry place, away from direct sunlight.

- Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

Off-white suspension with orange flavor.

- Ingredients

- Active ingredient

- Ibuprofen

- Inactive ingredients

- Potassium sorbate, Citric acid anhydrous, Trisodium citrate dihydrate, Saccharin sodium, Sodium chloride, Xanthan gum, Hypromellose, Maltitol syrup, Glycerin, Polysorbate 80, Orange flavor, Purified water.

- MAL number:

MALXXXXXXXXXX

Product Registration Holder

MEGA LIFESCIENCES SDN BHD
B-28-02, The Ascent, Paradigm,
No.1, Jalan SS7/26A, Kelana Jaya,
47301 Petaling Jaya, Selangor, Malaysia.

Manufacturer

MEGA LIFESCIENCES Public
Company Limited
515/1 Soi 8, Bangpoo Industrial Estate,
Pattana 3 Road, Moo 4, Phraeksa,
Mueang, Samutprakarn 10280, Thailand.

Date of revision

12/06/2026

Serial Number

NPRA (R2) 26/062