

THROZZ PARACETAMOL 500MG TABLET

Paracetamol (500mg)

What is in this leaflet

1. What THROZZ PARACETAMOL 500MG TABLET is used for
2. How THROZZ PARACETAMOL 500MG TABLET works
3. Before you use THROZZ PARACETAMOL 500MG TABLET
4. How to use THROZZ PARACETAMOL 500MG TABLET
5. While you are using it
6. Side effects
7. Storage and disposal of THROZZ PARACETAMOL 500MG TABLET
8. Product description
9. Manufacturer and Product registration holder
10. Date of revision
11. Serial number

What THROZZ PARACETAMOL 500MG TABLET is used for
THROZZ PARACETAMOL 500MG TABLET is used for relief of fever, pain and headache.

How THROZZ PARACETAMOL 500MG TABLET works

Paracetamol works by inhibiting the production of chemical compounds in the body called prostaglandin which causes pain and fever. This action helps to reduce pain and fever.

Before you use THROZZ PARACETAMOL 500MG TABLET

When you must not use it

Do not use THROZZ PARACETAMOL 500MG TABLET - if you are allergic (hypersensitive) to paracetamol or any of the other ingredients of this medicine.

Before you start to use it

- Do not take more than recommended dose as it may cause serious harm to your liver or kidney.
- This medicine contains paracetamol. Do not take any other paracetamol containing medicines at the same time.

Talk to your doctor or pharmacist before taking THROZZ

PARACETAMOL 500MG TABLET if:

- you have liver or kidney problems.
- you have glutathione depleted or are a chronic heavy alcohol user. It may increase the risk of metabolic acidosis. Signs of metabolic acidosis include: deep, rapid, difficult breathing, nausea, vomiting, loss of appetite. Contact a doctor immediately if you get a combination of these symptoms.
- you have non-serious arthritis that requires painkiller every day.
- you are pregnant, planning for pregnancy or breast-feeding.
- Please see your doctor if your symptoms do not improve.

Taking other medicines

- Tell your doctor if you are taking warfarin or blood thinning agent. Taking paracetamol and warfarin for a prolonged period may increase risk of bleeding.
- Tell your doctor if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket, or health food shop.

How to use THROZZ PARACETAMOL 500MG TABLET

How much to take

- Adults (including the elderly) and children aged 12 years and above: 1 - 2 tablets.

Maximum daily dose: 4000mg (8 tablets).

- Children, 6 to 11 years: ½ - 1 tablet.

Maximum daily dose: 2000mg (4 tablets).

- Do not give to children under the age of 12 years old.
- Oral administration only.

When to use it

To be taken every 4-6 hours as required or as directed by your doctor or pharmacist.

How long to use it

Do not take for a long period of time. Refer to your pharmacist or doctor for advice.

Please see your doctor if your

symptoms do not improve.

If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

If you use too much (overdose)

Contact your doctor immediately or go to the emergency department of the nearest hospital if you find yourself or anyone else may have used too much of this medicine. Do this even there are no signs of discomfort or poisoning because of the risk of liver failure. You may need urgent medical attention.

While you are using it

Things you must do

- Take the medicine exactly as your doctor has told you or according to instruction on pack.
- Tell all the doctors, dentists and pharmacists treating you that you are taking THROZZ PARACETAMOL 500MG TABLET
- Tell your doctor immediately if you become pregnant while taking this medicine.

Things you must not do

- Do not stop using the medicine unless advised by your doctor.
- Do not exceed the stated dose.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give THROZZ PARACETAMOL 500MG TABLET to anyone else even if they have the same symptoms or condition as you.

Things to be careful of

ALLERGY ALERT:

Paracetamol may cause severe skin reactions. Symptoms may include skin reddening, blisters, and rash. These could be signs of a serious condition. If these reactions occur, stop use and seek medical assistance right away.

Driving and using machines

THROZZ PARACETAMOL 500MG TABLET

Paracetamol (500mg)

THROZZ PARACETAMOL 500MG TABLET is unlikely to cause an effect ability to drive or use machines.

Side Effects

Stop taking this medicine and tell your doctor right away if you have any of the following side effects:

- You experience allergic reactions such as skin rash or itching, sometimes accompanied by breathing problems, or swelling of the lips, tongue, throat, or face.
- Experience a skin peeling, or mouth ulcers.
- Experience respiratory problems with aspirin or non-steroidal anti-inflammatory drugs, and experiences a similar reaction to this product.
- Experience unexplained bruising or bleeding.

These reactions are rare.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of THROZZ PARACETAMOL 500MG TABLET

Storage

Store below 30°C.

Keep this medicine out of the sight and reach of children.

Disposal

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

Product Description

What it looks like

White, capsule shaped tablet. Plain on one side with a break line on the other side.

Ingredients

Active Ingredient

Paracetamol 500mg

Inactive ingredients

Maize Starch

Pregelatinized Maize Starch

Povidone

Sodium Starch Glycolate

Magnesium Stearate

Purified Water

MAL number

MAL xxxxxxxxxxxx

Manufacturer

Steril-Gene Life Sciences (P) Ltd
No.45, Mangalam Main Road,
Mangalam Village, Villianur
Commune,
Puducherry – 605110, India

Product Registration Holder

Medicinspire Global Sdn Bhd
B-5-28, Ativo Plaza
No.1, Jalan PJU 9/1
Damansara Avenue
Bandar Sri Damansara
52200 Kuala Lumpur
Malaysia

Date of revision

04/01/2026

Serial Number

NPRA (R2) 25/238