

THROZZ BROMHEXINE 8MG TABLETS

Bromhexine Hydrochloride (8mg)

What is in this leaflet

1. What THROZZ BROMHEXINE 8MG TABLETS is used for
2. How THROZZ BROMHEXINE 8MG TABLETS works
3. Before you use THROZZ BROMHEXINE 8MG TABLETS
4. How to use THROZZ BROMHEXINE 8MG TABLETS
5. While you are using it
6. Side effects
7. Storage and disposal of THROZZ BROMHEXINE 8MG TABLETS
8. Product description
9. Manufacturer and Product registration holder
10. Date of revision

What THROZZ BROMHEXINE

8MG TABLETS is used for THROZZ BROMHEXINE 8MG TABLETS is used for the reduction of sputum viscosity.

How THROZZ BROMHEXINE 8MG TABLETS works

THROZZ BROMHEXINE 8MG TABLETS is a mucolytic (agent which dissolve thick mucus) expectorant whose action is to stimulate the mucous glands to produce a secretion and reduced content of acid glycoprotein fibres.

Before you use THROZZ BROMHEXINE 8MG TABLETS

When you must not use it

Do not use THROZZ BROMHEXINE 8MG TABLETS if you are allergic (hypersensitive) to Bromhexine Hydrochloride.

Before you start to use it

Tell your doctor or pharmacist if:

- you are allergic to any other medicines.
- you are taking any other medicines including those not prescribed by

- your doctor.
- you are taking any dietary supplements.
- you have gastric ulceration.

Taking other medicines

Seek for your doctor or pharmacist's advice if you intend to use THROZZ

Bromhexine 8mg Tablet together with other medication. You must also inform them about all the medicines you are taking, including:
- Erythromycin, cephalixin and amoxicillin

How to use THROZZ BROMHEXINE 8MG TABLETS

How much to take

Adults:
1-2 tablets (8-16mg) 3-4 times daily

Children 5-10 years: ½ tablet (4mg), 3 times daily

Children under 5 years:
½ tablet (4mg), 2 times daily

When to use it

3 times daily.

How long to use it

THROZZ BROMHEXINE 8MG TABLETS should be taken as your doctor advised you to do so. Do not stop taking this medicine without talking to the doctor. Please see your doctor if your symptoms do not improve.

If you forget to use it

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the medicine and skip the missed dose. Do not take a double dose to make up for the missed dose.

If you use too much (overdose)

Contact your doctor immediately or go to the emergency department of the nearest hospital if you find yourself or anyone else may have used too much of this medicine. Do this even there are no signs of discomfort or poisoning.

You may need urgent medical attention.

While you are using it

Things you must do

- Take the medicine exactly as your doctor has told you or according to instruction on pack.
- Tell all the doctors, dentists and pharmacists treating you that you are taking THROZZ BROMHEXINE 8MG TABLETS

Things you must not do

- Do not stop using the medicine unless advised by your doctor.
- Do not exceed the stated dose.
- Do not take any new medicines without consulting your doctor or pharmacist.
- Do not give THROZZ BROMHEXINE 8MG TABLETS to anyone else even if they have the same symptoms or condition as you.

Things to be careful of

- Take with caution for patients with gastric ulceration.
- Do not take THROZZ BROMHEXINE 8MG TABLETS during the first trimester of pregnancy and when breastfeeding.

Side Effects

All medicines can cause unwanted side effects although not everyone experience it. See a doctor immediately if you have any of the following serious side effects. You may need urgent medical treatment.

- Stomach and intestinal disturbances.
- Transitory rise in serum transaminase (An enzyme that is normally present in liver and heart cells, and is released into blood when the liver or heart is damaged) levels.

THROZZ BROMHEXINE 8MG TABLETS may cause severe allergy and serious skin reactions. Stop using this medicine and seek medical assistance immediately if you experience any of the following symptoms:

THROZZ BROMHEXINE 8MG TABLETS

Bromhexine Hydrochloride (8mg)

- Severe allergy: breathing difficulties, light headedness, skin swellings or rash.
- Severe skin reaction: skin reddening, blisters, rash, fever, sore throat or eye irritation.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Product Registration Holder

Medicinspire Global Sdn Bhd B-5-28,
Ativo Plaza
No.1, Jalan PJU 9/1 Damansara
Avenue Bandar Sri Damansara 52200
Kuala Lumpur Malaysia

Date of revision

27/03/2026

Serial Number

NPRA (R2) 26/041

Storage and Disposal of THROZZ BROMHEXINE 8MG TABLETS

Storage

Store below 30°C. Keep this medicine out of the sight and reach of children.

Disposal

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

Product Description

What it looks like

White to off white, round tablet, bevel-edged on one side and break-line on other side.

Ingredients

Active Ingredient

Bromhexine Hydrochloride 8mg

Inactive ingredients

Lactose Monohydrate 200M

Maize Starch

Magnesium Stearate

MAL number

MAL26056033XZ

Manufacturer

SAGA LIFESCIENCES LIMITED

Survey No. 198/2 & 198/3

Chachrawadi Vasna Ta.Sanand

District Ahmedabad

382210 India