

CAVSAMIR

Sevelamer carbonate Tablets 800 mg

What is in this leaflet

1. What Cavsamir is used for
2. How Cavsamir works
3. Before you use Cavsamir
4. How to use Cavsamir
5. While you are using it
6. Side effects
7. Storage and Disposal of Cavsamir
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision
11. Serial number

What Cavsamir is used for

Cavsamir is used for patients who have kidneys that do not work properly are not able to control the level of serum phosphorus in their blood. The amount of phosphate then rises (your doctor will call this hyperphosphataemia).

Increased levels of serum phosphorus can lead to hard deposits in your body called calcification. These deposits can stiffen your blood vessels and make it harder for blood to be pumped around the body. Increased serum phosphorus can also lead to itchy skin, red eyes, bone pain and fractures.

How Cavsamir works

Cavsamir contains sevelamer carbonate as the active ingredient. It binds phosphate from food in the digestive tract and so reduces serum phosphorus levels in the blood.

Before you use Cavsamir

-When you must not use it

Do not take Cavsamir if:

- You are allergic (hypersensitive) to sevelamer carbonate or to any of the other ingredients of this medicine (listed in Section 8 below).
- Signs of an allergic reaction include: a rash, swallowing or breathing problems, swelling of your lips, face, throat or tongue
- You have low levels of phosphate in your blood (your doctor will check this for you)
- You have bowel obstruction.

-Before you start to use it

Check with your doctor before taking Cavsamir if:

- You have swallowing problems
- You have problems related with motility (movement) in your stomach

and bowel

- You are being sick frequently
- You have active inflammation of the bowel
- You have undergone major surgery on your stomach or bowel
- You have severe constipation

Additional treatments:

Due to either your kidney condition or your dialysis treatment you may:

- Develop low or high levels of calcium in your blood. Since Cavsamir does not contain calcium your doctor might prescribe additional calcium tablets.
- Have a low amount of vitamin D in your blood. Therefore, your doctor may monitor the levels of vitamin D in your blood and prescribe additional vitamin D as necessary. If you do not take multivitamin supplements you may also develop low levels of vitamins A, E, K and folic acid in your blood and therefore your doctor may monitor these levels and prescribe supplemental vitamins as necessary.

Taking other medicines

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Cavsamir should not be taken at the same time as ciprofloxacin (an antibiotic).

If you are taking medicines for heart rhythm problems or for epilepsy, you should consult your doctor when taking Cavsamir.

The effects of medicines such as ciclosporin, mycophenolate mofetil and tacrolimus (medicines used to suppress the immune system) may be reduced by Cavsamir.

Your doctor will advise you if you are taking these medicines. Thyroid hormone deficiency may uncommonly be observed in certain people taking levothyroxine (used to treatment low thyroid hormone levels) and Cavsamir. Therefore your doctor may monitor the levels of thyroid stimulating hormone in your blood more closely.

How to use Cavsamir

Always take Cavsamir exactly as your doctor has told you. You should check with your doctor or pharmacist if you are not sure.

- Swallow your tablets orally in whole with plenty of water.
- Do not crush, chew or break into pieces.
- You must take Cavsamir with meals.

How much to use

Your doctor will prescribe Cavsamir to you based on your serum phosphorus level.

The recommended starting dose of Cavsamir tablets for adults and the elderly (> 65 years) is one to two tablets of 800mg with each meal, 3 times a day.

In some cases where Cavsamir should be taken at the same time as another medicine. Your doctor may advise you to take this medicine 1 hour before or 3 hours after Cavsamir intake, or they may consider monitoring the blood levels of that medicine.

Cavsamir is not recommended for use in children and adolescents below 18 years of age.

-When to use it

Take Cavsamir at about the same time each day unless your doctor tells you otherwise.

-How long to use it

Continue taking your medicine for as long as your doctor tells you.

-If you forget to use it

- If you forget to take your tablet, leave out that dose completely and take the next tablet at the usual time.
- Do not take a double dose to make up for a forgotten tablet.
- If you have any further questions on the use of this product, ask your doctor or pharmacist.

-If you use too much (overdose)

There are no reported overdoses in patients. In the event of a possible overdose you should contact your doctor

CAVSAMIR

Sevelamer carbonate Tablets 800 mg

immediately. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

While you are using it

-Things you must do

Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking any medicine if:

- You are pregnant, think you may be pregnant or are planning to have a baby. It is unknown whether Cavsamir has any effect on unborn babies.
- You wish to breast-feed your baby. It is unknown whether Cavsamir may pass through breast milk and affect your baby.

-Things you must not do

- Do not give this medicine to anyone else, even if they have the same condition as you.
- Do not use this medicine to treat any other complaints unless your doctor tells you to.
- Do not stop taking Cavsamir, or change the dosage, without checking with your doctor.

Things to be careful of

Driving and using machines

No studies on the effects on the ability to drive and use machines has been performed. If you are affected, do not drive and do not use any tools or machines.

Side Effect

Like all medicines, Cavsamir can cause side effects, although not everyone gets them.

Do not be alarmed by this list of possible side effects. You may not experience any of them. Tell your doctor immediately

Very common side effects

- Constipation
- Vomiting
- Upper abdominal pain
- Nausea

Common side effects

- Diarrhoea
- Abdominal pain
- Indigestion
- Flatulence

Cases of hypersensitivity, itching,

rash, slow intestine motility (movement)/blockages in the intestine, and perforation in the intestine wall have been reported.

Since constipation may be an early symptom of a blockage in your intestine, please inform your doctor or pharmacist.

Other side effects not listed above may occur in some people.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of Cavsamir

If you have any queries about any aspect of your medicine, or any questions regarding the information in this leaflet, discuss them with your doctor or pharmacist.

Storage

Store below 30 Celcius. Dispense in a tight container. Protect from moisture. Keep out of reach and sight of children.

Disposal

Medicines should not be disposed of via wastewater or household waste. Ask your pharmacist how to dispose of medicines no longer required. These measures will help to protect the environment.

Product Description

-What it looks like

White to off-white, oval, film coated tablets plain on one side and imprinted 'R 789' on other side.

Ingredients

- Active Ingredient

Sevelamer Carbonate

- Inactive Ingredients

Mannitol, Crospovidone Type B (XL-10), Hydroxypropyl Cellulose (Klucel LF), Colloidal Silicon Dioxide, Zinc Stearate, Opadry AMB Translucent (OY-B-29000), Purified water, Opacode Black S-1-17823

MAL number:

MAL21066027AZ

Manufacturer

Dr. Reddy's Laboratories Limited,
FTO SEZ - Process Unit 1,
Survey No.'s 57 to 59, 60, 62 & 72,
Sector No.9 to 14 & 17 to 20,
Devunipalavalasa (V), Ranasthalam (M),
Srikakulam District - 532409,
Andhra Pradesh, India.

Product Registration Holder

GoodScience Sdn Bhd
No.7, Jalan KPK 4/3,
Kawasan Perindustrian Kundang,
Kundang Jaya, 48020 Rawang,
Selangor, Malaysia

Date of revision

09/06/2025

Serial Number

NPRA(R2) 20/116