Medinazole Plus Cream in all its dosages is indicated for the treatment of dermatomycoses complicated by inflammatory and/or pruritic manifestations of skin disorders.

Directions of Use: For topical use only. Medinazole Plus Cream should be applied to the affected area no more than 2 times daily, preferably once in the morning and once in the evening. Medinazole Plus Cream should not be applied with an occlusive dressing, or to large skin areas of the body.

Precautions: For external use only. Avoid in contact with eyes, nose and mouth. Not to be taken. If symptoms persist, please consult a doctor.

Storage: Store below 30°C. Protect from light. Keep cap tightly closed. Keep out of reach of children. Jauhkan daripada capaian kanak-kanak.

Please refer enclosed leaflet for more information.

15g

Medicare+

Medinazole Plus Cream

Econazole Nitrate (1%w/w) Triamcinolone Acetonide (0.1%w/w)

Active Ingredients:
Econazole Nitrate 1%w/w
Triamcinolone Acetonide 0.1%w/w
Preservative: Chlorocresol 0.1%w/w

Product Registration Holder
WORLD MEDICARE SUPPLIES SDN BHD
PT 5740(1st Floor), Jalan TS 2/1E,
Taman Semarak 2, 71800 Nilai,
Negeri Sembilan, Malaysia.

Controlled Medicine

Manufacturer
WINWA MEDICAL SDN. BHD.
1952, Lorong I.K.S. Bukit Minyak 3,
Taman I.K.S. Bukit Minyak,
14000 Bukit Mertajam,
Pulau Pinang, Malaysia.

Registration No.: MAL00000000AZC

9 555838 TEMP

A.EWmNP15-B.01

Medinazole Plus Cream

BN :
MFG :
EXP :

Size: 120mm x 25mm x 35mm

C M Y K

Medinazole - Outer Box
Option B