

THROATSIL LOZENGES (MENTHOL FLAVOUR)

2,4-Dichlorobenzyl Alcohol 1.2 mg;

Amylmetacresol 0.6 mg

In this leaflet:

1. What THROATSIL LOZENGES is used for?
2. How THROATSIL LOZENGES works?
3. Before you use THROATSIL LOZENGES
4. How to use THROATSIL LOZENGES?
5. While you are using THROATSIL LOZENGES
6. Side effects
7. Storage and Disposal of THROATSIL LOZENGES
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What THROATSIL LOZENGES is used for?

THROATSIL LOZENGES contains 2,4-Dichlorobenzyl Alcohol and Amylmetacresol used for relief of mouth and throat infections for adults and children. These ingredients have antiseptic property which kill the bacteria associated with mouth and throat infections. By dissolving lozenge slowly in the mouth allows the active ingredients to work in the discomfort and soothe the painful area.

How THROATSIL LOZENGES works

THROATSIL LOZENGES helps to relieve the soreness and discomfort of mouth and throat infections. It contains the antiseptic ingredients, such as 2,4-Dichlorobenzyl Alcohol and Amylmetacresol which kill the bacteria that cause the mouth and throat infection. It is available in the form of lozenge which slowly dissolve in the mouth when sucked.

Before you use THROATSIL LOZENGES

- When you must not use it

Do not take this product if you are allergic (hypersensitive) to 2,4-Dichlorobenzyl Alcohol, Amylmetacresol or any of the other inactive ingredients

Pregnancy and breast-feeding

Ask your doctor or pharmacist for advice before taking THROATSIL LOZENGES.

- Before you start to use it

Tell your doctor or pharmacist if you:

- Are taking or have recently taken any other medicines, including medicines obtained without a prescription.
- Suffer or are suffering from any other mouth and throat problems.

- Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, including any that you buy without a prescription from a pharmacy, supermarket or healthcare shop.

How to use THROATSIL LOZENGES

- How much to use

Place lozenges into the mouth allowing it to dissolve slowly. Dissolve 1 - 3 lozenges in mouth slowly in a divided dose per day.

Remember that young children can choke on lozenges.

- When to use it

Use THROATSIL LOZENGES when necessary or as directed by your doctor or pharmacist.

- How long to use it

Continue taking THROATSIL LOZENGES as long as it is necessary or as recommended by your doctor or pharmacist.

- If you forget to use it

Take THROATSIL LOZENGES as soon as possible or consult your doctor or pharmacist on what you should do if you forget to use it.

Take the missed dose as soon as you remember. If it is almost time for your next dose, wait until then to take the lozenge and skip the missed dose. Do not take a double dose to make up for the missed dose.

- If you use too much (overdose)

If you take more of THROATSIL LOZENGES than you should, you may experience stomach (gastrointestinal) discomfort.

Do not take any more of THROATSIL LOZENGES and consult your doctor or pharmacist.

Contact your doctor immediately or go to the Emergency Department of your nearest hospital, if you think you or anyone else may have taken too much of THROATSIL LOZENGES. You may need urgent medical attention even if there is no sign of discomfort or poisoning.

While you are using THROATSIL LOZENGES

- Things you must do

Take your medicine exactly as your doctor has told you.

THROATSIL LOZENGES (MENTHOL FLAVOUR)

2,4-Dichlorobenzyl Alcohol 1.2 mg;
Amylmetacresol 0.6 mg

Tell all the doctors, dentists and pharmacists treating you that you are taking THROATSIL LOZENGES.

Tell your doctor immediately if you become pregnant while taking THROATSIL LOZENGES

- Things you must not do

Do not stop taking the medicine unless advised by your doctor or pharmacist.

Do not take any new medicines without consulting your doctor or pharmacist.

Do not give THROATSIL LOZENGES to anyone else, even if they have the same symptoms or condition as you.

Side effects

Like all medicines, THROATSIL LOZENGES may cause side effects, although not everybody gets them. Allergic (hypersensitivity) reactions may be occurred when you are using it.

Visit your doctor or pharmacist immediately if you experience any side effects after taking this medicine.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by calling Tel: 03-78835550, or visiting the website portal.npra.moh.gov.my (Public → Reporting Medicinal Problems / Side Effects / AEFI).

Storage and Disposal of THROATSIL LOZENGES

- Storage

Keep out of the reach of children.

Store below 30°C.

- Disposal

THROATSIL LOZENGES should not be disposed of via wastewater or household waste. Ask your doctor or pharmacist on how to dispose it when it is no longer required. These measures will help to protect the environment.

Product Description

- What it looks like

A blue round biconvex lozenge.
Plain on both sides.

- Ingredients

- Active ingredients

2,4-Dichlorobenzyl Alcohol 1.2 mg;
Amylmetacresol 0.6 mg

- Inactive ingredients

Sucrose, Liquid Glucose, Brilliant Blue FCF (CI. No.42090), Menthol, Eucalyptus Oil, Citric Acid.

Registration No.

MAL*****

Manufacturer:

Millimed Co., Ltd.
193 Moo 1, Suksawad Road,
Pak Khlong Bang Plakot,
Phra Samut Chedi,
Samut Prakan 10290, Thailand

Product Registration Holder:

HNB Alliance SDN. BHD.
No. 6, Jalan Industri Mas 3,
Taman Mas Sepang,
47100 Puchong,
Selangor, Malaysia.

Date of revision:

10/12/2017