

VOMIZ 8 (Ondansetron Tablet USP 8mg)

Ondansetron 8mg (as hydrochloride dihydrate)

What is in this leaflet

1. What VOMIZ 8 is used for
2. How VOMIZ 8 works
3. Before you use VOMIZ 8
4. How to use VOMIZ 8
5. While you are using it
6. Side effects
7. Storage and Disposal of VOMIZ 8
8. Product Description
9. Manufacturer and Product Registration Holder
10. Date of revision

What VOMIZ 8 is used for

It is used to prevent and manage nausea (feeling sick) and vomiting (being sick), which can occur:

- during treatment for cancer (chemotherapy or radiotherapy)
- after an operation

How VOMIZ 8 works

VOMIZ 8 belongs to a group of medicines called anti-emetics. Ondansetron is a 5HT₃ receptor antagonist. It works by inhibiting the 5HT₃ receptors on neurons located in peripheral and central nervous system of the body.

Before you use VOMIZ 8

When you must not use it

Do not take VOMIZ 8 if you are

- if you are taking apomorphine (used to treat Parkinson's disease, a movement disorder)
- if you are allergic (hypersensitive) to ondansetron or any other ingredients of VOMIZ 8

If you think this applies to you, do not use VOMIZ 8 until you have checked with your doctor.

Before you start to use it

Before you use VOMIZ 8 your doctor needs to know:

- if you are allergic to medicines similar to VOMIZ 8, such as medicines containing granisetron or palonosetron
- if you have ever had heart problems, such as an irregular heartbeat (arrhythmia)
- if you have bowel problems
- if you have liver disease, your doctor may lower your dose of VOMIZ 8
- if you are taking other medicines (interactions with other medicinal products including vaccines or biologics)

Check with your doctor if you think any of these may apply to you.

Pregnancy and breast-feeding

VOMIZ 8 is not recommended for use during pregnancy.

- Tell your doctor if you are pregnant or planning to become pregnant. VOMIZ 8 may harm your unborn baby.
- If you do become pregnant during treatment with VOMIZ 8, tell your doctor.

Breast-feeding is not recommended during treatment with VOMIZ 8. The ingredients can pass into your breast milk, and may affect your baby. Talk to your doctor about this.

Females of child-bearing potential and male patients

If you are a woman of childbearing age, your doctor will check if you are pregnant and perform a pregnancy test if necessary before starting treatment with VOMIZ 8. If you may become pregnant, you should use effective birth control during treatment and for at least 2 days after stopping VOMIZ 8.

Ask your doctor about options of effective birth control.

Taking other medicines

Tell your doctor or pharmacist if you are taking any other medicines, if you have taken any recently, or if you start taking new ones. This includes medicines bought without a prescription.

- carbamazepine or phenytoin (used to treat epilepsy)
- rifampicin (used to treat infections such as tuberculosis (TB), type of bacterial infection)
- fluoxetine, paroxetine, sertraline, fluvoxamine, citalopram, escitalopram (SSRIs [selective serotonin reuptake inhibitors]) (used to treat depression and/or anxiety)
- venlafaxine, duloxetine (SNRIs [serotonin noradrenaline reuptake inhibitors]) (used to treat depression and/or anxiety)
- tramadol (a pain killer)

How to use VOMIZ 8

How much to use

Always take VOMIZ 8 exactly as your doctor has told you to. Check with your doctor or pharmacist if you are not sure.

To prevent you from feeling sick and vomiting after cancer treatment.

On the day of treatment:

- The usual dose of VOMIZ 8 is 8 mg to be taken one to two hours before treatment, followed by 8 mg every 12 hours.

On the following days:

Adults

- The usual dose is 8 mg twice a day, for up to 5 days
- Children and Adolescents (ages 3 years to 17 years)
- Your doctor will decide on the correct dose of VOMIZ 8 for your child. The usual dose is up to 8 mg a day, for up to 5 days.

To prevent you from feeling sick and vomiting after an operation

Adults

- The usual adult dose is 16 mg one hour before your operation. Children and Adolescents (ages 3 years to 17 years)
- VOMIZ 8 tablets are not recommended for use in children after an operation

When to use it

Use as directed by your doctor or pharmacist.

How long to use it

Take VOMIZ 8 for as long as your doctor recommends. Do not stop unless your doctor advises you to.

If you forget to use it

Take your dose as soon as possible then take your next dose at the usual time. Do not take a double dose to make up for a missed dose.

If you are not sure what to do, ask your doctor or pharmacist.

If you use too much (overdose)

If you accidentally use too much VOMIZ 8, it is unlikely to cause you any problems. But if you are worried, or you feel unwell, ask your doctor or pharmacist for advice.

While you are using it

Things you must do

VOMIZ 8 (Ondansetron Tablet USP 8mg)

Ondansetron 8mg (as hydrochloride dihydrate)

Check with your doctor or pharmacist before taking your medicine and take it as prescribed by your doctor.

Swallow the tablets whole with a glass of water.

Tell your doctor or nurse if you continue to feel sick.

Things you must not do

This medicine has been prescribed for you. Do not pass it on to others. It may harm them, even if their symptoms are the same as yours.

Things to be careful of

Tell your doctor or pharmacist immediately if you get any of these symptoms during and after the treatment with VOMIZ 8.

- if you experience sudden chest pain or chest tightness (myocardial ischemia).

VOMIZ 8 tablets are not recommended for use in children and Adolescents (aged 3 years to 17 years) after an operation.

Side effects

Like all medicines, VOMIZ 8 can cause side effects, although not everybody gets them.

Some side effects could be serious

STOP taking or receiving VOMIZ 8 and seek medical help immediately if you experience any of the following:

Severe allergic reactions

These are rare in people taking VOMIZ 8. Signs include:

- raised and itchy rash (hives)
- swelling, sometimes of the face or mouth (angioedema), causing difficulty in breathing
- collapse

Myocardial ischemia: Signs include:

- sudden chest pain or
- chest tightness

Other possible side effects

Other side effects include the following listed below. If these side effects become severe, please tell your doctor, pharmacist or healthcare provider.

Very common: may affect more than 1 in 10 people

- headache

Common: may affect up to 1 in 10 people

- feeling of warmth or flushing
- constipation

Uncommon: may affect up to 1 in 100 people

- fits (seizures)
- involuntary muscle movements or spasms
- irregular or slow heartbeat
- chest pain
- low blood pressure
- hiccups
- increases in substances (enzymes) produced by the liver (may show up in blood test).

Rare: may affect up to 1 in 1,000 people

- severe allergic reactions
- disturbance in heart rhythm (sometimes causing a sudden loss of consciousness)
- dizziness
- temporary blurred or altered vision

Very rare side: may affect up to 1 in 10,000 people:

- a widespread rash with blisters and skin peeling on much of the body surface (toxic epidermal necrolysis)
- temporary blindness

If you notice any side effects not listed in this leaflet, please inform your doctor or pharmacist.

You may report any side effects or adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my [Consumers → Reporting Side Effects to Medicines (ConSERF) or Vaccines (AEFI)].

Storage and Disposal of VOMIZ 8

Storage

Store below 30°C.

Keep out of the sight and reach of children.

Do not use your medicine after the expiry date which is stated on the pack after 'EXP'. The expiry date refers to the last day of that month.

Disposal

Do not dispose of medicines in your wastewater or household rubbish. Ask your pharmacist how to dispose of medicines no longer required. This will help to protect the environment.

Product Description

What it looks like

Yellow coloured, capsule shaped, biconvex, film coated tablets plain on both the sides.

Ingredients

Active ingredient

Each tablet contains Ondansetron (as hydrochloride dihydrate) 8mg

Inactive ingredient

Microcrystalline Cellulose, Lactose Anhydrous, Pregelatinised Starch, Magnesium stearate, Opadry Yellow 03B82493 (HPMC 2910/ HYPROMELLOSE 6cP, TITANIUM DIOXIDE, MACROGOUPEG 400, IRON OXIDE YELLOW), Purified water

Nature and Contents of Container

VOMIZ 8 is available in blister packs of 30 tablets.

MAL number:

MAL24066019AZ

Manufacturer

Zydus Lifesciences Limited, Kundaim Industrial Estate, Plot No. 203-213, Kundaim, Goa - 403 115, INDIA.

Product Registration Holder

Mansa Healthcare Sdn Bhd
B-05-3A, 3 Two Square, No 2, Jalan 19/1, 46300 Petaling Jaya, Selangor Malaysia

Date of revision

1-Dec-2024

Serial Number

NPRA (R2) 23/078