

PREFER 60mg

Pyridostigmine Bromide Tablet 60mg)

What is in this leaflet

1. What PREFER is used for
2. How PREFER works
3. Before you use PREFER
4. How to use PREFER
5. While you are taking PREFER
6. Side effects
7. Storage and Disposal of PREFER
8. Product Description
9. Manufacturer
10. Product Registration holder
11. Serial Number

What PREFER is used for

PREFER is used for the treatment of myasthenia gravis. This is a condition whereby you have muscle weakness and tiredness especially on repeated use.

How PREFER works

PREFER belongs to a group of medicines called cholinergic antimusclerics. It works by preventing the breakdown of a chemical called acetylcholine. This chemical is needed to stimulate muscle movement.

Before you use PREFER

- When you must not use it

Do not take PREFER if you are:

- allergic (hypersensitive) to pyridostigmine, bromides or any of the other ingredients of PREFER.
- suffering from mechanical obstruction (blockage) of the intestinal or urinary tract.

If you are not sure whether you should start taking this medicine, talk to your doctor.

- Before you start to use it

Tell your doctor if you have had any unusual or allergic reactions to anticholinesterase agents. Other substances such as foods, preservatives or dyes. Symptoms of an allergic reaction may include shortness of breath, wheezing or difficulty

breathing, swelling of the face, lips, tongue or other parts of the body, rash, itching or hives on the skin.

- Taking other medicines

Tell your doctor if you have or have had any of the following medical conditions (ie: kidney problem) or taking any other medicines, including medicines obtained without a prescription.

Some medicines and PREFER may interfere with each other. These include:

- muscle relaxants
- atropine contained in medicines for travel sickness and stomach cramps, or use in surgery (injection) or used as an antidote to insecticide poisoning (tablets)

Your doctor or pharmacist may have more information on medicines to be careful with or to avoid while taking medication.

- Pregnancy and Lactation

The safety of PREFER for use during pregnancy has not been established. Hence, it should only be used strictly as directed by your doctor.

Since it is not known whether the active ingredient in PREFER passes into breast milk, nursing mothers are advised to refrain from breast feeding their babies during treatment.

Your doctor can advise you of the risks and benefits of taking PREFER while you are pregnant or breast feeding.

Ask your doctor or pharmacist for advice before taking any medicine.

How to use PREFER

- How much to use

When using PREFER, it is important to remember that the full effect appears gradually, usually within 15-30 minutes. You should try to take

your medicine so that it can work when your muscles are needed most, for example when getting up and at meal times.

Swallow the tablets whole with water. If you have difficulty swallowing, you can break the tablets into pieces.

The number of tablets you take depends on your illness and age. Also, the number of doses you take each day, the interval between doses should be at least 6 hours.

Adult: The usual adult dose is 1 to 3 tablets, 2 to 4 times daily or higher doses if required.

Children: For children under 6 years the usual dose is half a tablet (30mg). For children aged 6 to 12 years, the usual starting dose is 1 tablet (60mg). The doctor will then gradually increase the dose until maximum improvement is seen. In children the total dose per day is usually 30 to 360mg. Follow all directions given to you by your doctor and pharmacist carefully. They may differ from the information contained in this leaflet. If you do not understand the instructions on the label, ask your doctor or pharmacist for help.

- When to take it

PREFER should be taken 30 to 60 minutes before meals.

- How long to take it

It is important that you keep taking these tablets until your doctor tells you to stop.

- If you forget to take it

If you forget to take a dose, take it as soon as you remember. Do not take a double dose to make up for a forgotten dose, just carry on as before.

- If you take too much (overdose)

If you take too many tablets or someone else accidentally takes your

PREFER 60mg

Pyridostigmine Bromide Tablet 60mg)

medicine, contact your doctor, pharmacist or nearest hospital straight away.

Taking too many tablets may cause pronounced muscle weakness or exacerbated muscle weakness, bradycardia (slow heart rate), tachycardia (fast heart rate)

While you are taking PREFER

- Things you must do

If you are about to be started on any new medicine, remind your doctor and pharmacist that you are taking PREFER.

Tell any other doctors, dentists and pharmacists who treat you that you are taking this medicine.

If you are going to have surgery, tell the surgeon or anaesthetist that you are taking this medicine. It may affect other medicines used during surgery.

If you become pregnant while taking this medicine, tell your doctor immediately. If you are about to have any blood test, tell your doctor that you are taking this medicine. It may interfere with the results of some tests.

- Things you must not do

Do not take PREFER to treat any other complaints unless your doctor tells you to.

Do not give your medicine to anyone else, even if they have the same condition as you.

- Things to be careful of

Do not take the medicine if the packaging appears to have been tampered with or if the medicine shows any signs of damage.

You must not take the medicine after the expiry date shown on the packaging.

- Driving and using machines

The effects of PREFER on the ability to drive and operate machinery have not been established. PREFER contains lactose. Therefore, if you have been told by your doctor that you have intolerance to some sugars, contact your doctor before taking this medicinal product.

Side effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while you are taking PREFER. You may need medical attention if you get some of the side effects.

Do not be alarmed by the following lists of side effects. You may not experience any of them.

Tell your doctor or pharmacist if you notice any of the following and they worry you:

- nausea and vomiting
- diarrhea
- stomach cramps or pains
- increased saliva
- skin rash

The above list includes serious side effects which may require urgent medical attention. Tell your doctor or pharmacist if you notice anything else that is making you feel unwell. Other side effects not listed above may also occur in some people.

You may report any side effects of adverse drug reactions directly to the National Centre for Adverse Drug Reaction Monitoring by visiting the website npra.gov.my (Consumer → Reporting → Reporting Side Effects to Medicine (ConSERF) or Vaccines (AEFI))

Storage and Disposal of PREFER

- Storage

Store below 30°C. Store in the original package to protect from light and moisture.

- Disposal

You must not take the medicine after the expiry date shown on the packaging. If you have any unused medicine, return it to your pharmacist who will dispose of it safely.

Product Description

- What it looks like

White to off white, round, flat-faced tablet debossed with "C60" one side and quadrisect scored on one side and plain on the other side.

- Ingredients

Active ingredients:

- Pyridostigmine Bromide

Inactive ingredients:

Anhydrous Lactose, Colloidal Silicon Dioxide, Stearic Acid

MAL number:

MALxxxxxxxxAZ

Manufacturer

Centaur Pharmaceuticals Pvt. Ltd.
Plot No.4, Hinjewadi Phase II,
International Biotech Park, Pune,
Maharashtra, IN-411057 India

Product Registration Holder:

Mansa Healthcare Sdn Bhd
B-05-3A, 3 Two Square, No 2, Jalan
19/1, 46300 Petaling Jaya, Selangor,
Malaysia

Date of revision

1-Mar-2025

Serial Number

NPRA (R2) 24/105